

RLBP1 Antibody (monoclonal) (M02)**Mouse monoclonal antibody raised against a full length recombinant RLBP1.****Catalog # AT3650a****Specification**

RLBP1 Antibody (monoclonal) (M02) - Product Information

Application	IF, WB, E
Primary Accession	P12271
Other Accession	BC004199
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 kappa
Calculated MW	36474

RLBP1 Antibody (monoclonal) (M02) - Additional Information**Gene ID** 6017**Other Names**

Retinaldehyde-binding protein 1, Cellular retinaldehyde-binding protein, RLBP1, CRALBP

Target/Specificity

RLBP1 (AAH04199, 1 a.a. ~ 317 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

RLBP1 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


RLBP1 Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


RLBP1 Antibody (monoclonal) (M02) - Images




Immunofluorescence of monoclonal antibody to RLBP1 on HeLa cell. [antibody concentration 20 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (60.61 KDa) .



RLBP1 monoclonal antibody (M02), clone 4H5 Western Blot analysis of RLBP1 expression in HepG2 (Cat # AT3650a)



Detection limit for recombinant GST tagged RLBP1 is approximately 0.1ng/ml as a capture antibody.

RLBP1 Antibody (monoclonal) (M02) - Background

The protein encoded by this gene is a 36-kD water-soluble protein which carries 11-cis-retinaldehyde or 11-cis-retinal as physiologic ligands. It may be a functional component of the visual cycle. Mutations of this gene have been associated with severe rod-cone dystrophy, Bothnia dystrophy (nonsyndromic autosomal recessive retinitis pigmentosa) and retinitis punctata albescens.

RLBP1 Antibody (monoclonal) (M02) - References

Central retinal findings in Bothnia dystrophy caused by RLBP1 sequence variation. Burstedt MS, et al. Arch Ophthalmol, 2010 Aug. PMID 20696998. Development of a Diagnostic Genetic Test for Simplex and Autosomal Recessive Retinitis Pigmentosa. Clark GR, et al. Ophthalmology, 2010 Jun 28. PMID 20591486. Bothnia dystrophy is caused by domino-like rearrangements in cellular retinaldehyde-binding protein mutant R234W. He X, et al. Proc Natl Acad Sci U S A, 2009 Nov 3. PMID 19846785. Prefrontal cortex shotgun proteome analysis reveals altered calcium homeostasis and immune system imbalance in schizophrenia. Martins-de-Souza D, et al. Eur Arch Psychiatry Clin Neurosci, 2009 Apr. PMID 19165527. Carrier of R14W in carbonic anhydrase IV presents Bothnia dystrophy phenotype caused by two allelic mutations in RLBP1. Kohn L, et al. Invest Ophthalmol Vis Sci, 2008 Jul. PMID 18344446.