

RNF2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant RNF2. Catalog # AT3672a

Specification

RNF2 Antibody (monoclonal) (M01) - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Calculated MW

WB, E
O99496
NM_007212
Human, Mouse
mouse
Monoclonal
IgG2a Kappa
37655

RNF2 Antibody (monoclonal) (M01) - Additional Information

Gene ID 6045

Other Names

E3 ubiquitin-protein ligase RING2, 632-, Huntingtin-interacting protein 2-interacting protein 3, HIP2-interacting protein 3, Protein DinG, RING finger protein 1B, RING1b, RING finger protein 2, RING finger protein BAP-1, RNF2, BAP1, DING, HIPI3, RING1B

Target/Specificity

RNF2 (NP_009143, 192 a.a. \sim 290 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

RNF2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

RNF2 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

• Western Blot




- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

RNF2 Antibody (monoclonal) (M01) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .




RNF2 monoclonal antibody (M01), clone 6C2. Western Blot analysis of RNF2 expression in Raw 264.7 ((Cat # AT3672a)






RNF2 monoclonal antibody (M01), clone 6C2 Western Blot analysis of RNF2 expression in NIH/3T3 ((Cat # AT3672a)



Western Blot analysis of RNF2 expression in transfected 293T cell line by RNF2 monoclonal antibody (M01), clone 6C2.

Lane 1: RNF2 transfected lysate(37.7 KDa).


Lane 2: Non-transfected lysate.



Western blot analysis of RNF2 over-expressed 293 cell line, cotransfected with RNF2 Validated



Chimera RNAi ((Cat # AT3672a)



Detection limit for recombinant GST tagged RNF2 is approximately 0.03ng/ml as a capture antibody.

RNF2 Antibody (monoclonal) (M01) - Background

Polycomb group (PcG) of proteins form the multiprotein complexes that are important for the transcription repression of various genes involved in development and cell proliferation. The protein encoded by this gene is one of the PcG proteins. It has been shown to interact with, and suppress the activity of, transcription factor CP2 (TFCP2/CP2). Studies of the mouse counterpart suggested the involvement of this gene in the specification of anterior-posterior axis, as well as in cell proliferation in early development. This protein was also found to interact with huntingtin interacting protein 2 (HIP2), an ubiquitin-conjugating enzyme, and possess ubiquitin ligase activity.

RNF2 Antibody (monoclonal) (M01) - References

1.Scmh1 Has E3 Ubiquitin Ligase Activity for Geminin and Histone H2A and Regulates Geminin Stability Directly or Indirectly via Transcriptional Repression of Hoxa9 and Hoxb4.Yasunaga S, Ohtsubo M, Ohno Y, Saeki K, Kurogi T, Tanaka-Okamoto M, Ishizaki H, Shirai M, Mihara K, Brock HW, Miyoshi J, Takihara Y.Mol Cell Biol. 2013 Feb;33(4):644-60. doi: 10.1128/MCB.00974-12.2.Inaugural Article: Distinct histone modifications in stem cell lines and tissue lineages from the early mouse embryo.Rugg-Gunn PJ, Cox BJ, Ralston A, Rossant J.Proc Natl Acad Sci U S A. 2010 May 17. [Epub ahead of print]3.Polycomb-group complex 1 acts as an E3 ubiquitin ligase for Geminin to sustain hematopoietic stem cell activity.Ohtsubo M, Yasunaga S, Ohno Y, Tsumura M, Okada S, Ishikawa N, Shirao K, Kikuchi A, Nishitani H, Kobayashi M, Takihara Y.Proc Natl Acad Sci U S A. 2008 Jul 29;105(30):10396-401. Epub 2008 Jul 23.