

SCAND1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant SCAND1. Catalog # AT3778a

Specification

SCAND1 Antibody (monoclonal) (M01) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW

WB, IF, E <u>P57086</u> <u>BC036709</u> Human mouse Monoclonal IgG2b Kappa 19082

SCAND1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 51282

Other Names SCAN domain-containing protein 1, SCAND1, SDP1

Target/Specificity SCAND1 (AAH36709, 1 a.a. ~ 179 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 IF~~1:50~200 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions SCAND1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

SCAND1 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides



- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

SCAND1 Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to SCAND1 on HeLa cell . [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (45.43 KDa) .



Detection limit for recombinant GST tagged SCAND1 is approximately 0.1ng/ml as a capture antibody.

SCAND1 Antibody (monoclonal) (M01) - Background



The SCAN domain is a highly conserved, leucine-rich motif of approximately 60 aa originally found within a subfamily of zinc finger proteins. This gene belongs to a family of genes that encode an isolated SCAN domain, but no zinc finger motif. Functional studies have established that the SCAN box is a protein interaction domain that mediates both hetero- and homoprotein associations, and maybe involved in regulation of transcriptional activity. Two transcript variants with different 5' UTRs, but encoding the same protein, have been described for this gene.

SCAND1 Antibody (monoclonal) (M01) - References

Multiple genetic variants along candidate pathways influence plasma high-density lipoprotein cholesterol concentrations. Lu Y, et al. J Lipid Res, 2008 Dec. PMID 18660489.Spectroscopic characterization of the tumor antigen NY-REN-21 and identification of heterodimer formation with SCAND1. Carneiro FR, et al. Biochem Biophys Res Commun, 2006 Apr 28. PMID 16540086.The LIFEdb database in 2006. Mehrle A, et al. Nucleic Acids Res, 2006 Jan 1. PMID 16381901.Towards a proteome-scale map of the human protein-protein interaction network. Rual JF, et al. Nature, 2005 Oct 20. PMID 16189514.From ORFeome to biology: a functional genomics pipeline. Wiemann S, et al. Genome Res, 2004 Oct. PMID 15489336.