

STK32C Antibody (monoclonal) (M02)**Mouse monoclonal antibody raised against a full length recombinant STK32C.****Catalog # AT4080a****Specification**

STK32C Antibody (monoclonal) (M02) - Product Information

Application	WB, IHC
Primary Accession	Q86UX6
Other Accession	BC015792
Reactivity	Human, Mouse
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	54994

STK32C Antibody (monoclonal) (M02) - Additional Information**Gene ID** 282974**Other Names**Serine/threonine-protein kinase 32C, PKE, Yet another novel kinase 3, STK32C
{ECO:0000312|EMBL:AAH457601}, YANK3**Target/Specificity**

STK32C (AAH15792, 1 a.a. ~ 369 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

STK32C Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


STK32C Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


STK32C Antibody (monoclonal) (M02) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (66.33 kDa) .



STK32C monoclonal antibody (M02), clone 3B4 Western Blot analysis of STK32C expression in NIH/3T3 (Cat # AT4080a)



Immunoperoxidase of monoclonal antibody to STK32C on formalin-fixed paraffin-embedded

human prostate. [antibody concentration 3 ug/ml]

STK32C Antibody (monoclonal) (M02) - Background

The protein encoded by this gene is a member of the serine/threonine protein kinase family. The specific function of this kinase is not known. [provided by RefSeq]