

**TAF1 Antibody (monoclonal) (M01)****Mouse monoclonal antibody raised against a partial recombinant TAF1.****Catalog # AT4136a****Specification**

---

**TAF1 Antibody (monoclonal) (M01) - Product Information**

Application	WB, E
Primary Accession	<a href="#">P21675</a>
Other Accession	<a href="#">NM_004606</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	214714

**TAF1 Antibody (monoclonal) (M01) - Additional Information****Gene ID** 6872**Other Names**

Transcription initiation factor TFIID subunit 1, Cell cycle gene 1 protein, TBP-associated factor 250 kDa, p250, Transcription initiation factor TFIID 250 kDa subunit, TAF(II)250, TAFII-250, TAFII250, TAF1, BA2R, CCG1, CCGS, TAF2A

**Target/Specificity**

TAF1 (NP\_004597, 1784 a.a. ~ 1893 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

E~~N/A

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

TAF1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**TAF1 Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## TAF1 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .



Detection limit for recombinant GST tagged TAF1 is approximately 0.1ng/ml as a capture antibody.

## TAF1 Antibody (monoclonal) (M01) - Background

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is the basal transcription factor TFIID, which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes the largest subunit of TFIID. This subunit binds to core promoter sequences encompassing the transcription start site. It also binds to activators and other transcriptional regulators, and these interactions affect the rate of transcription initiation. This subunit contains two independent protein kinase domains at the N and C-terminals, but also possesses acetyltransferase activity and can act as a ubiquitin-activating/conjugating enzyme. This gene is part of a complex transcriptional unit

(TAF1/DYT3), wherein some products share exons with TAF1 as well as additional exons downstream.

#### **TAF1 Antibody (monoclonal) (M01) - References**

TAF1 differentially enhances androgen receptor transcriptional activity via its N-terminal kinase and ubiquitin-activating and -conjugating domains. Tavassoli P, et al. Mol Endocrinol, 2010 Apr. PMID 20181722. TAF1 interacts with and modulates human papillomavirus 16 E2-dependent transcriptional regulation. Centeno F, et al. Intervirology, 2008. PMID 18580066. Transcription factor TAFII250 phosphorylates the acidic domain of Mdm2 through recruitment of protein kinase CK2. Allende-Vega N, et al. Mol Cell Biochem, 2008 Sep. PMID 18548200. Search for cellular partners of human papillomavirus type 16 E2 protein. Olejnik-Schmidt AK, et al. Arch Virol, 2008. PMID 18305892. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.