

THEA Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant THEA.****Catalog # AT4232a****Specification**

THEA Antibody (monoclonal) (M01) - Product Information

| | |
|-------------------|--------------------------|
| Application | WB, E |
| Primary Accession | Q8WXI4 |
| Other Accession | BC001517 |
| Reactivity | Human |
| Host | mouse |
| Clonality | Monoclonal |
| Isotype | IgG1 Kappa |
| Calculated MW | 68492 |

THEA Antibody (monoclonal) (M01) - Additional Information**Gene ID** 26027**Other Names**

Acyl-coenzyme A thioesterase 11, Acyl-CoA thioesterase 11, 312-, Acyl-CoA thioester hydrolase 11, Adipose-associated thioesterase, Brown fat-inducible thioesterase, BFIT, ACOT11, BFIT, KIAA0707, THEA

Target/Specificity

THEA (AAH01517, 1 a.a. ~ 110 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

THEA Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


THEA Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

THEA Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 KDa) .



Detection limit for recombinant GST tagged ACOT11 is approximately 0.03ng/ml as a capture antibody.

THEA Antibody (monoclonal) (M01) - Background

This gene encodes a protein with acyl-CoA thioesterase activity towards medium (C12) and long-chain (C18) fatty acyl-CoA substrates. Expression of a similar murine protein in brown adipose tissue is induced by cold exposure and repressed by warmth. Expression of the mouse protein has been associated with obesity, with higher expression found in obesity-resistant mice compared with obesity-prone mice. Alternative splicing results in two transcript variants encoding different isoforms.

THEA Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Gene-centric association signals for lipids and apolipoproteins identified via

the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. A revised nomenclature for mammalian acyl-CoA thioesterases/hydrolases. Hunt MC, et al. J Lipid Res, 2005 Sep. PMID 16103133. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.