

**TRIM37 Antibody (monoclonal) (M01)****Mouse monoclonal antibody raised against a partial recombinant TRIM37.****Catalog # AT4355a****Specification**

---

**TRIM37 Antibody (monoclonal) (M01) - Product Information**

Application	WB, E
Primary Accession	<a href="#">O94972</a>
Other Accession	<a href="#">NM_015294</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	107906

**TRIM37 Antibody (monoclonal) (M01) - Additional Information****Gene ID** 4591**Other Names**

E3 ubiquitin-protein ligase TRIM37, 632-, Mulibrey nanism protein, Tripartite motif-containing protein 37, TRIM37, KIAA0898, MUL, POB1

**Target/Specificity**

TRIM37 (NP\_056109, 865 a.a. ~ 964 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

E~~N/A

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

TRIM37 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**TRIM37 Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TRIM37 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



Detection limit for recombinant GST tagged TRIM37 is approximately 0.3ng/ml as a capture antibody.

### TRIM37 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the tripartite motif (TRIM) family, whose members are involved in diverse cellular functions such as developmental patterning and oncogenesis. The TRIM motif includes zinc-binding domains, a RING finger region, a B-box motif and a coiled-coil domain. The RING finger and B-box domains chelate zinc and might be involved in protein-protein and/or protein-nucleic acid interactions. The gene mutations are associated with mulibrey (muscle-liver-brain-eye) nanism, an autosomal recessive disorder that involves several tissues of mesodermal origin. Alternatively spliced transcript variants encoding the same protein have been identified.

### TRIM37 Antibody (monoclonal) (M01) - References

Shifted Transversal Design smart-pooling for high coverage interactome mapping. Xin X, et al. Genome Res, 2009 Jul. PMID 19447967. Gynecological tumors in Mulibrey nanism and role for RING finger protein TRIM37 in the pathogenesis of ovarian fibrothecomas. Karlberg S, et al. Mod Pathol, 2009 Apr. PMID 19329943. A novel mutation in TRIM37 is associated with mulibrey nanism in a

Turkish boy. Do?anc T, et al. Clin Dysmorphol, 2007 Jul. PMID 17551331. Wilms' tumor and novel TRIM37 mutations in an Australian patient with mulibrey nanism. H?m?l?inen RH, et al. Clin Genet, 2006 Dec. PMID 17100991. Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. Olsen JV, et al. Cell, 2006 Nov 3. PMID 17081983.