

ZNF274 Antibody (monoclonal) (M04)**Mouse monoclonal antibody raised against a partial recombinant ZNF274.****Catalog # AT4621a****Specification**

ZNF274 Antibody (monoclonal) (M04) - Product Information

Application	WB, IF
Primary Accession	O96GC6
Other Accession	NM_133502
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	74177

ZNF274 Antibody (monoclonal) (M04) - Additional Information**Gene ID** 10782**Other Names**

Neurotrophin receptor-interacting factor homolog, Zinc finger protein 274, Zinc finger protein HFB101, Zinc finger protein with KRAB and SCAN domains 19, Zinc finger protein zfp2, Zf2, ZNF274, ZKSCAN19

Target/Specificity

ZNF274 (NP_598009, 420 a.a. ~ 530 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IF~~1:50~200

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

ZNF274 Antibody (monoclonal) (M04) is for research use only and not for use in diagnostic or therapeutic procedures.


ZNF274 Antibody (monoclonal) (M04) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF274 Antibody (monoclonal) (M04) - Images




Immunofluorescence of monoclonal antibody to ZNF274 on HeLa cell . [antibody concentration 10 ug/ml]



Western Blot analysis of ZNF274 expression in transfected 293T cell line by ZNF274 monoclonal antibody (M04), clone 1D8.

Lane 1: ZNF274 transfected lysate(74.2 KDa).

Lane 2: Non-transfected lysate.



Western blot analysis of ZNF274 over-expressed 293 cell line, cotransfected with ZNF274 Validated Chimera RNAi (Cat # AT4621a)

ZNF274 Antibody (monoclonal) (M04) - Background

This gene encodes a zinc finger protein containing five C2H2-type zinc finger domains, one or two Kruppel-associated box A (KRAB A) domains, and a leucine-rich domain. The encoded protein has been suggested to be a transcriptional repressor. It localizes predominantly to the nucleolus. Alternatively spliced transcript variants encoding different isoforms exist. These variants utilize alternative polyadenylation signals.

ZNF274 Antibody (monoclonal) (M04) - References

Systematic identification of SH3 domain-mediated human protein-protein interactions by peptide array target screening. Wu C, et al. *Proteomics*, 2007 Jun. PMID 17474147. The KAP1 corepressor functions to coordinate the assembly of de novo HP1-demarcated microenvironments of heterochromatin required for KRAB zinc finger protein-mediated transcriptional repression. Sripathy SP, et al. *Mol Cell Biol*, 2006 Nov. PMID 16954381. Large-scale cDNA transfection screening for genes related to cancer development and progression. Wan D, et al. *Proc Natl Acad Sci U S A*, 2004 Nov 2. PMID 15498874. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. *Genome Res*, 2004 Oct. PMID 15489334. Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. *Nat Genet*, 2004 Jan. PMID 14702039.