

APC Anti-Mouse CD11c (N418) Antibody
Catalog # ATB10013**Specification**

APC Anti-Mouse CD11c (N418) Antibody - Product Information

Application	FC
Isotype	Armenian Hamster IgG
Concentration	0.2 mg/mL
Reactivity	Mouse
Formulation	10 mM NaH ₂ PO ₄ , 150 mM NaCl, 0.09% Na ₃ N, 0.1% gelatin, pH7.2
Host	Armenian Hamster

APC Anti-Mouse CD11c (N418) Antibody - Additional Information

Gene ID	16411
Gene Name	Itgax
Alternative Name(s)	
p150, integrin α X, Itgax, CR4	

Format
APC**Preparation**

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Application Notes

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.

Storage Conditions

2-8°C protected from light


APC Anti-Mouse CD11c (N418) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

APC Anti-Mouse CD11c (N418) Antibody - Images



C57Bl/6 splenocytes were stained with Anti-Mouse MHC Class II FITC (35-5321) and 0.125 ug Anti-Mouse CD11c APC (ATB10013) (right panel) or 0.125 ug Armenian Hamster IgG APC (left panel).



C57Bl/6 splenocytes were stained with Anti-Mouse MHC Class II FITC and 0.125 ug APC Anti-Mouse CD11c (N418) manufactured by Tonbo Biosciences (left panel) or eBioscience (right panel).

APC Anti-Mouse CD11c (N418) Antibody - Background

The N418 antibody reacts with mouse CD11c, also known as integrin alpha X. This 150 kDa cell surface glycoprotein is part of a family of integrin receptors that mediate adhesion between cells (cell-cell) and components of the extracellular matrix, e.g. fibrinogen (cell-matrix). In addition, integrins are active signaling receptors which recruit leukocytes to inflammatory sites and promote cell activation. Complete, functional integrin receptors consist of distinct combinations of integrin chains which are differentially expressed. Integrin alpha X (CD11c) assembles with Integrin beta-2 (CD18) into a receptor complex known as CR4 which can bind and induce signaling through ICAMs and VCAM-1 on endothelial cells and can also facilitate removal of iC3b bearing foreign cells. The N418 antibody is widely used as a marker for CD11c expression on dendritic cells (DC), often in parallel with markers for CD11b, for identification of developmental stages and mature subsets of this cell type. CD11c is prominently expressed on tissue macrophages, and is also detected on some types of activated T cells and intestinal intraepithelial lymphocytes (IEL).