

**FITC Anti-Mouse CD127 (IL-7Ra) (A7R34) Antibody**  
**Catalog # ATB10121****Specification****FITC Anti-Mouse CD127 (IL-7Ra) (A7R34) Antibody - Product Information**

Application	FC
Isotype	Rat IgG2a, kappa
Concentration	0.5 mg/mL
Reactivity	Mouse
Formulation	10 mM NaH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, 0.09% NaN <sub>3</sub> , 0.1% gelatin, pH7.2
Host	Rat

**FITC Anti-Mouse CD127 (IL-7Ra) (A7R34) Antibody - Additional Information**

Gene ID	16172
Gene Name	Il17ra
<b>Alternative Name(s)</b>	

Interleukin-7 Receptor alpha, IL-7Ra

**Format**  
FITC**Preparation**

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

**Application Notes**

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.

**Storage Conditions**

2-8°C protected from light


**FITC Anti-Mouse CD127 (IL-7Ra) (A7R34) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

### **FITC Anti-Mouse CD127 (IL-7Ra) (A7R34) Antibody - Images**



C57Bl/6 splenocytes were stained with APC Anti-Mouse CD3 (20-0031) and 1 ug FITC Anti-Mouse CD127 (ATB10121) (right panel) or 1 ug FITC Rat IgG2a isotype control (left panel).

### **FITC Anti-Mouse CD127 (IL-7Ra) (A7R34) Antibody - Background**

The A7R34 antibody is specific for mouse CD127, a 60-90 kDa cell surface protein also known as the Interleukin-7 Receptor alpha chain, or IL-7R alpha. CD127 is typically expressed at the cell surface as a heterodimer with the common gamma chain (CD132). This complex acts as the functional receptor for IL-7, a cytokine important in T and B cell development, and in mature T cell homeostasis. A second cytokine known as Thymic Stromal Lymphopoietin (TSLP) also binds to a receptor complex of CD127 and the TSLPR chain to trigger activation of dendritic cells, and is involved in B cell development, allergy and autoimmunity. The A7R34 antibody may be used as a phenotypic marker for CD127 on immature B cells, on subsets of thymocytes which are double negative (CD4-CD8- ) or single positive (CD4+ or CD8+), and at low levels on mature, peripheral T cells. CD127 is a key marker, when used in combination with CD4 and CD25, to distinguish Treg and effector/memory Treg populations known as T(REM).