

In Vivo Ready[™] Anti-Human CD3 (OKT3) Antibody Catalog # ATB10139

Specification

In Vivo Ready[™] Anti-Human CD3 (OKT3) Antibody - Product Information

Application Isotype Concentration Reactivity Formulation Host IHC-P, IHC-F, FC, FA Mouse IgG2a, kappa 2 mg/mL Human 10 mM NaH2PO4, 150 mM NaCl, pH7.2 Mouse

In Vivo Ready[™] Anti-Human CD3 (OKT3) Antibody - Additional Information

Gene ID Gene Name Alternative Name(s) Leu-4, T3 916 CD3E

Format In Vivo Ready™

Preparation

This monoclonal antibody preparation was purified from tissue culture supernatant via affinity chromatography. For In Vivo Ready[™] (IVR) products, each preparation is also evaluated for endotoxin levels using the LAL assay. It is recommended to store the product undiluted at 4°C. Do not freeze.

Application Notes

This purified format is guaranteed to be >90% pure as determined by SDS-PAGE analysis. Citations are provided as a convenience to you - please consult Materials and Methods sections for additional details about the use of any product in these publications.

Endotoxin Level Less than or equal to 0.01 EU/ug, as determined by the LaL assay

Storage Conditions 2-8°C

In Vivo Ready[™] Anti-Human CD3 (OKT3) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence



- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

In Vivo Ready[™] Anti-Human CD3 (OKT3) Antibody - Images

In Vivo Ready[™] Anti-Human CD3 (OKT3) Antibody - Background

The OKT3 antibody is specific for human CD3e, also known as CD3 epsilon, a 20 kDa subunit of the T cell receptor complex along with CD3 gamma and CD3 delta. These integral membrane protein chains assemble with additional chains of the T cell receptor (TCR), as well as CD3 zeta chain, to form the T cell receptor - CD3 complex. Together with co-receptors CD4 or CD8, the complex serves to recognize antigens bound to MHC molecules on antigen-presenting cells. These interactions promote T cell receptor signaling (T cell activation), inducing a number of cellular responses including proliferation, differentiation, production of cytokines or activation-induced cell death. CD3 is differentially expressed during thymocyte-to-T cell development and on all mature T cells. The OKT3 antibody is a widely used phenotypic marker for human T cells. In addition, as the CD3e subunit of the TCR complex contains intracellular signaling domains, binding of OKT3 antibody to CD3e can induce cell activation. A recent publication of the crystal structure of a CD3e-mitogenic antibody complex provides further insight as to the action of commonly used agonist antibodies, as well as specific epitope-binding data for the widely used human CD3 antibodies OKT3 and UCHT1 (Fernandes, R.A. et al. 2012. J. Biol. Chem. 287: 13324-13335). OKT3 has also been shown to be cross-reactive with Chimpanzee CD3 and has been used for in vitro activation of T cells in this species.