

PE-Cy7 Anti-Human CD45RA (HI100) Antibody
Catalog # ATB10251**Specification****PE-Cy7 Anti-Human CD45RA (HI100) Antibody - Product Information**

| | |
|---------------|-------------------------------------------------------------|
| Application | FC |
| Isotype | Mouse IgG2b, kappa |
| Concentration | 5 uL (0.25 ug)/test |
| Reactivity | Human |
| Formulation | 10 mM NaH2PO4, 150 mM NaCl, 0.09% NaN3, 0.1% gelatin, pH7.2 |
| Host | Mouse |

PE-Cy7 Anti-Human CD45RA (HI100) Antibody - Additional Information

| | |
|-----------|-------|
| Gene ID | 5788 |
| Gene Name | PTPRC |
| Format | |
| PE-Cy7 | |

Preparation

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Application Notes

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5 uL per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 uL. The number of cells within a sample should be determined empirically, but typically ranges between 1x10e5 to 1x10e8 cells.


Storage Conditions

2-8°C protected from light

PE-Cy7 Anti-Human CD45RA (HI100) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PE-Cy7 Anti-Human CD45RA (HI100) Antibody - Images

Human peripheral blood lymphocytes were stained with 5 μ L (0.25 μ g) PE-Cy7 Anti-Human CD45RA (ATB10251) (solid line) or 0.25 μ g PE-Cy7 Mouse IgG2b isotype control (dashed line).

PE-Cy7 Anti-Human CD45RA (HI100) Antibody - Background

The HI100 antibody reacts with the human CD45 isoform known as CD45RA, a protein tyrosine phosphatase of 220 kDa. CD45 is one of the most abundant hematopoietic markers, and is expressed on all leukocytes (the Leukocyte Common Antigen, LCA). Various isoforms are generated and expressed in cell-specific patterns. With their broad cell distribution, CD45 isoforms are critical for many leukocyte functions, regulating signal transduction and cell activation associated with the T cell receptor, B cell receptor, and IL-2 receptor. Other forms of CD45, with restricted cellular expression, include CD45R (B220), CD45RB, CD45RO and others. The HI100 antibody is widely used as a marker for human CD45RA expression on naïve and activated T cells, B cells, and monocytes.