

**LYZ Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5031****Specification**

---

**LYZ Antibody (C-term) - Product Information**

Application	IHC-P, WB,E
Primary Accession	<a href="#">P61626</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=17 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**LYZ Antibody (C-term) - Additional Information****Gene ID** 4069**Antigen Region**  
119-154**Other Names**  
Lysozyme C, 4-beta-N-acetylmuramidase C, LYZ, LZM**Dilution**  
IHC-P~~1:25  
WB~~1:2000**Target/Specificity**  
This LYZ antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 119-154 amino acids from the C-terminal region of human LYZ.**Format**  
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.**Storage**  
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.**Precautions**  
LYZ Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.**LYZ Antibody (C-term) - Protein Information****Name** LYZ

## Synonyms LZM

## Function

Lysozymes have primarily a bacteriolytic function; those in tissues and body fluids are associated with the monocyte-macrophage system and enhance the activity of immunoagents.

## Cellular Location


Secreted.

## LYZ Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## LYZ Antibody (C-term) - Images




Western blot analysis of lysates from HepG2, THP-1, WiDr cell line and human spleen tissue lysate(from left to right), using LYZ Antibody (C-term)(Cat. #AW5031). AW5031 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.




All lanes : Anti-LYZ Antibody (C-term) at 1:2000 dilution Lane 1: WiDr whole cell lysate Lane 2: THP-1 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: Human spleen lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 17 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




All lanes : Anti-LYZ Antibody (C-term) at 1:2000 dilution Lane 1: THP-1 whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: Human spleen lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 15 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Immunohistochemical analysis of paraffin-embedded M. small intestine section using LYZ Antibody (C-term)(Cat#AW5031). AW5031 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.




Immunohistochemical analysis of paraffin-embedded R. small intestine section using LYZ Antibody (C-term)(Cat#AW5031). AW5031 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. small intestine section using LYZ Antibody (C-term)(Cat#AW5031). AW5031 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



AW5031 staining LYZ in human tonsil tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded Human spleen section using Pink1(Cat#AW5031). AW5031 was diluted at 1:1000 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

**LYZ Antibody (C-term) - Background**

Lysozymes have primarily a bacteriolytic function; those in tissues and body fluids are associated with the monocyte- macrophage system and enhance the activity of immunoagents.

**LYZ Antibody (C-term) - References**

Castanon M.J.,et al.Gene 66:223-234(1988).  
Chung L.P.,et al.Proc. Natl. Acad. Sci. U.S.A. 85:6227-6231(1988).  
Yoshimura K.,et al.Biochem. Biophys. Res. Commun. 150:794-801(1988).  
Peters C.W.B.,et al.Eur. J. Biochem. 182:507-516(1989).  
Huang B.,et al.Sheng Wu Hua Hsueh Tsa Chih 9:269-273(1993).