

USP5 Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AW5077

Specification

USP5 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW Isotype Antigen Source WB,E <u>P45974</u> Human, Mouse, Rat Mouse Monoclonal H=96;M=96;Rat=96 KDa IgG1,ĸ HUMAN

USP5 Antibody - Additional Information

Gene ID 8078

Antigen Region 1-160

Other Names

Ubiquitin carboxyl-terminal hydrolase 5, Deubiquitinating enzyme 5, Isopeptidase T, Ubiquitin thioesterase 5, Ubiquitin-specific-processing protease 5, USP5, ISOT

Dilution WB~~1:1000

Target/Specificity

This USP5 antibody is generated from a mouse immunized with a recombination protein from the human region of human USP5.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions USP5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

USP5 Antibody - Protein Information

Name USP5



Synonyms ISOT

Function

Deubiguitinating enzyme that participates in a wide range of cellular processes by specifically cleaving isopeptide bonds between ubiquitin and substrate proteins or ubiquitin itself. Affects thereby important cellular signaling pathways such as NF-kappa-B, Wnt/beta- catenin, and cytokine production by regulating ubiguitin-dependent protein degradation. Participates in the activation of the Wnt signaling pathway by promoting FOXM1 deubiquitination and stabilization that induces the recruitment of beta-catenin to Wnt target gene promoter (PubMed:26912724). Regulates the assembly and disassembly of heat-induced stress granules by mediating the hydrolysis of unanchored ubiquitin chains (PubMed:29567855). Promotes lipopolysaccharide-induced apoptosis and inflammatory response by stabilizing the TXNIP protein (PubMed:37534934). Affects T-cell biology by stabilizing the inhibitory receptor on T-cells PDC1 (PubMed: 37208329). Acts as a negative regulator of autophagy by regulating ULK1 at both protein and mRNA levels (PubMed:37607937). Acts also as a negative regulator of type I interferon production by simultaneously removing both 'Lys-48'-linked unanchored and 'Lys-63'-linked anchored polyubiguitin chains on the transcription factor IRF3 (PubMed:39761299). Modulates the stability of DNA mismatch repair protein MLH1 and counteracts the effect of the ubiquitin ligase UBR4 (PubMed: 39032648). Upon activation by insulin, it gets phosphorylated through mTORC1-mediated phosphorylation to enhance YTHDF1 stability by removing 'Lys-11'-linked polyubiquitination (PubMed: 39900921). May also deubiquitinate other substrates such as the calcium channel CACNA1H (By similarity).

Cellular Location Cytoplasm. Cytoplasm, Stress granule. Nucleus

USP5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- **USP5 Antibody Images**





Western blot analysis of lysates from mouse brain tissue,PC-3 cell line,rat liver tissue,U-251 MG cell line,rat brain tissue (from left to right), using USP5 Antibody(Cat. #AW5077). AW5077 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysates at 20ug per lane.

USP5 Antibody - Background

Cleaves linear and branched multiubiquitin polymers with a marked preference for branched polymers. Involved in unanchored 'Lys-48'-linked polyubiquitin disassembly. Binds linear and 'Lys- 63'-linked polyubiquitin with a lower affinity. Knock-down of USP5 causes the accumulation of p53/TP53 and an increase in p53/TP53 transcriptional activity because the unanchored polyubiquitin that accumulates is able to compete with ubiquitinated p53/TP53 but not with MDM2 for proteasomal recognition.

USP5 Antibody - References

Falquet L.,et al.FEBS Lett. 376:233-237(1995). Ansari-Lari M.A.,et al.Genome Res. 6:314-326(1996). Ansari-Lari M.A.,et al.Genome Res. 7:268-280(1997). Tashayev V.L.,et al.Submitted (NOV-1995) to the EMBL/GenBank/DDBJ databases. Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.