

**APG5L (ATG5) Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AW5079**

**Specification**

---

**APG5L (ATG5) Antibody - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q9H1Y0</a>
Other Accession	<a href="#">Q3MQ04</a> , <a href="#">Q3MQ24</a>
Reactivity	Human, Mouse, Rat
Predicted	Pig, Bovine
Host	Mouse
Clonality	Monoclonal
Calculated MW	H=32,23;M=32;Rat=25 KDa
Isotype	IgG1, $\kappa$
Antigen Source	HUMAN

**APG5L (ATG5) Antibody - Additional Information**

**Gene ID** 9474

**Other Names**

Autophagy protein 5, APG5-like, Apoptosis-specific protein, ATG5, APG5L, ASP

**Dilution**

WB~~1:1000

**Target/Specificity**

This ATG5 antibody is generated from a mouse immunized with a recombination protein from the human region of human ATG5.

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

APG5L (ATG5) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**APG5L (ATG5) Antibody - Protein Information**

**Name** ATG5 ([HGNC:589](#))

**Synonyms** APG5L, ASP

**Function**

Involved in autophagic vesicle formation. Conjugation with ATG12, through a ubiquitin-like conjugating system involving ATG7 as an E1-like activating enzyme and ATG10 as an E2-like conjugating enzyme, is essential for its function. The ATG12-ATG5 conjugate acts as an E3-like enzyme which is required for lipidation of ATG8 family proteins and their association to the vesicle membranes. Involved in mitochondrial quality control after oxidative damage, and in subsequent cellular longevity. Plays a critical role in multiple aspects of lymphocyte development and is essential for both B and T lymphocyte survival and proliferation. Required for optimal processing and presentation of antigens for MHC II. Involved in the maintenance of axon morphology and membrane structures, as well as in normal adipocyte differentiation. Promotes primary ciliogenesis through removal of OFD1 from centriolar satellites and degradation of IFT20 via the autophagic pathway. As part of the ATG8 conjugation system with ATG12 and ATG16L1, required for recruitment of LRRK2 to stressed lysosomes and induction of LRRK2 kinase activity in response to lysosomal stress (By similarity).

**Cellular Location**

Cytoplasm. Preautophagosomal structure membrane; Peripheral membrane protein.  
Note=Colocalizes with nonmuscle actin. The conjugate detaches from the membrane immediately before or after autophagosome formation is completed (By similarity). Also localizes to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme. Under starved conditions, the ATG12-ATG5-ATG16L1 complex is translocated to phagophores driven by RAB33B (PubMed:32960676). {ECO:0000250, ECO:0000269|PubMed:32960676}

**Tissue Location**


Ubiquitous. The mRNA is present at similar levels in viable and apoptotic cells, whereas the protein is dramatically highly expressed in apoptotic cells

**APG5L (ATG5) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**APG5L (ATG5) Antibody - Images**



Western blot analysis of lysates from rat C6, HeLa, mouse NIH/3T3, SH-SY5Y, U-87 MG cell line (from left to right), using ATG5 Antibody (Cat. #AW5079). AW5079 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody.

#### **APG5L (ATG5) Antibody - Background**

Involved in autophagy vesicles formation. Conjugation with ATG12 through an ubiquitin-like conjugating system involving ATG7 as an E1-like activating enzyme and ATG10 as an E2-like conjugating enzyme, is essential for its function. The ATG12-ATG5 conjugate acts as an E3-like enzyme which is required for lipidation of ATG8 family proteins and their association to the vesicle membranes. Involved in mitochondrial quality control after oxidative damage, and in subsequent cellular longevity. The ATG12- ATG5 conjugate also regulates negatively the innate antiviral immune response by blocking the type I IFN production pathway through direct association with RARRES3 and MAVS. Plays also a role in translation or delivery of incoming viral RNA to the translation apparatus. HCV utilizes ATG5 as a proviral factor during the onset of viral infection. Plays a critical role in multiple aspects of lymphocyte development and is essential for both B and T lymphocyte survival and proliferation. Required for optimal processing and presentation of antigens for MHC II. Involved in the maintenance of axon morphology and membrane structures; as well as in normal adipocyte differentiation. Promotes primary ciliogenesis through removal of OFD1 from centriolar satellites and degradation of IFT20 via the autophagic pathway.

#### **APG5L (ATG5) Antibody - References**

Hammond E.M., et al. FEBS Lett. 425:391-395(1998).  
Chen Y., et al. Submitted (AUG-2000) to the EMBL/GenBank/DDBJ databases.  
Bechtel S., et al. BMC Genomics 8:399-399(2007).  
Mungall A.J., et al. Nature 425:805-811(2003).  
Grand R.J.A., et al. Exp. Cell Res. 218:439-451(1995).