

# **CABLES2 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5114

## **Specification**

## **CABLES2 Antibody (Center) - Product Information**

Application WB,E
Primary Accession Q9BTV7
Other Accession Q8K3M5

Reactivity Human, Mouse Host Rabbit

Clonality
Calculated MW
Polyclonal
H=52;M=53 KDa

Isotype Rabbit IgG
Antigen Source Human

## **CABLES2 Antibody (Center) - Additional Information**

**Gene ID 81928** 

**Antigen Region** 

182-210

#### **Other Names**

CABLES2;C20orf150; CDK5 and ABL1 enzyme substrate 2; CDK5 and ABL1 enzyme substrate 2; Interactor with CDK3 2

## **Dilution**

WB~~1:1000

### **Target/Specificity**

This CABLES2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 182-210 amino acids from the Central region of human CABLES2.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at  $2-8^{\circ}$ C for up to 2 weeks. For long term storage store at  $-20^{\circ}$ C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

CABLES2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **CABLES2 Antibody (Center) - Protein Information**



#### Name CABLES2

### Synonyms C20orf150

#### **Function**


Unknown. Probably involved in G1-S cell cycle transition.

# **CABLES2 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **CABLES2 Antibody (Center) - Images**



Western blot analysis of lysates from PC-3 cell line and mouse testis tissue(from left to right), using CABLES2Antibody(Center)(Cat. #AW5114). AW5114 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysates at 20ug per lane.

## CABLES2 Antibody (Center) - Background

Unknown. Probably involved in G1-S cell cycle transition.

## **CABLES2 Antibody (Center) - References**

Deloukas P., et al. Nature 414:865-871(2001). Daub H., et al. Mol. Cell 31:438-448(2008). Oppermann F.S., et al. Mol. Cell. Proteomics 8:1751-1764(2009).