

ESR2 Antibody(C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5284

Specification

ESR2 Antibody(C-term) - Product Information

Application WB,E
Primary Accession Q92731

Other Accession Q9XSB5, NP 001035365.1, Q9TU15

Reactivity Human

Predicted Bovine, Sheep
Host Rabbit
Clonality Polyclonal
Calculated MW H=55 KDa
Isotype Rabbit IgG

Antigen Source HUMAN

ESR2 Antibody(C-term) - Additional Information

Gene ID 2100

Antigen Region

451-480

Other Names

ESR2; ESTRB; NR3A2; Estrogen receptor beta; Nuclear receptor subfamily 3 group A member 2

Dilution

WB~~1:1000

Target/Specificity

This ESR2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 451-480 amino acids from the C-terminal region of human ESR2.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ESR2 Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ESR2 Antibody(C-term) - Protein Information



Name ESR2

Synonyms ESTRB, NR3A2

Function

Nuclear hormone receptor. Binds estrogens with an affinity similar to that of ESR1/ER-alpha, and activates expression of reporter genes containing estrogen response elements (ERE) in an estrogen- dependent manner (PubMed:20074560).

Cellular Location

 $Nucleus \ \{ ECO: 0000255 | PROSITE-ProRule: PRU00407, \ ECO: 0000269 | PubMed: 19126643, \ ECO: 0000269 | PubMed: 20074560 \}$

Tissue Location


[Isoform 1]: Expressed in testis and ovary, and at a lower level in heart, brain, placenta, liver, skeletal muscle, spleen, thymus, prostate, colon, bone marrow, mammary gland and uterus Also found in uterine bone, breast, and ovarian tumor cell lines, but not in colon and liver tumors. [Isoform 4]: Expressed in the testis. [Isoform 6]: Expressed in testis, placenta, skeletal muscle, spleen and leukocytes, and at a lower level in heart, lung, liver, kidney, pancreas, thymus, prostate, colon, small intestine, bone marrow, mammary gland and uterus. Not expressed in brain.

ESR2 Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ESR2 Antibody(C-term) - Images





Tel: 858.875.1900 Fax: 858.875.1999

Western blot analysis of lysates from MCF-7 cell line and human planceta tissue (from left to right), using ESR2 Antibody (C-term)(Cat. #AW5284). AW5284 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

ESR2 Antibody(C-term) - Background

This gene encodes a member of the family of estrogen receptors and superfamily of nuclear receptor transcription factors. The gene product contains an N-terminal DNA binding domain and C-terminal ligand binding domain and is localized to the nucleus, cytoplasm, and mitochondria. Upon binding to 17beta-estradiol or related ligands, the encoded protein forms homo- or hetero-dimers that interact with specific DNA sequences to activate transcription. Some isoforms dominantly inhibit the activity of other estrogen receptor family members. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been fully characterized.

ESR2 Antibody(C-term) - References

Fang, Y.J., et al. Tumour Biol. 31(6):651-658(2010) Hayes, D.F., et al. Clin. Pharmacol. Ther. 88(5):626-629(2010) Romero, R., et al. Am. J. Obstet. Gynecol. 203 (4), 361 (2010): Wurster, M., et al. Oncol. Rep. 24(3):653-659(2010) Paulus, J.K., et al. Lung Cancer (2010) In press: