

# Anti-Glargine Insulin polyclonal antibody

Purified Guinea pig polyclonal Antibody (Pab) Catalog # AW5686

### **Specification**

### Anti-Glargine Insulin polyclonal antibody - Product Information

Application Host Clonality **Antigen Source** 

sELISA,E **Guinea Pig Polyclonal** 

Insulin glargine, marketed under the names Lantus, among others, is a long-acting basal insulin analogue, given once daily to help control the blood sugar level of those with diabetes. Insulin glargine has a substitution of glycine for asparagine at N21 (Asn21) and two arginines added to the carboxy terminal of B chain. The arginine amino acids shift the isoelectric point from a pH of 5.4 to 6.7, making the molecule more soluble at an acidic pH and less soluble at physiological pH. The isoelectric shift also allows for the subcutaneous injection of a clear solution. The glycine substitution prevents deamidation of the acid-sensitive asparagine at acidic pH. In the neutral subcutaneous space, higher-order aggregates form, resulting in a slow, peakless dissolution and absorption of insulin from the site of injection. It can achieve a peakless level for at least 24 hours. Molecular formula: C267H404N72O78S6 Molecular weight:6063

### Anti-Glargine Insulin polyclonal antibody - Additional Information

**Other Names** Anti-Glargine Insulin polyclonal antibody

Dilution sELISA~~N/A


Target/Specificity

Guinea pig polyclonal antibody raised against Glargine Insulin .

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a antigen affinity purification column, followed by dialysis against PBS.







**Storage** 

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

Anti-Glargine Insulin polyclonal antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Anti-Glargine Insulin polyclonal antibody - Protein Information

### Anti-Glargine Insulin polyclonal antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Anti-Glargine Insulin polyclonal antibody - Images




Plate was coated with Glargine Insulin at 1.25 µg/ml in PBS, and then incubated with Anti-Glargine Insulin polyclonal antibody from 0.001 μg/ml to 0.5 μg/ml. The secondary antibody, HRP conjugated goat anti-Guinea pig IgG, were used at 1:6000 dilution.