

Zebrafish znf143 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb18703c**Specification**

Zebrafish znf143 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q1LYE3
Reactivity	Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	564-596

Zebrafish znf143 Antibody (C-term) - Additional Information**Gene ID** 393954**Other Names**

Zinc finger protein 143, Selenocysteine tRNA gene transcription-activating factor, znf143, staf

Target/Specificity

This Zebrafish znf143 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 564-596 amino acids of zebrafish znf143.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Zebrafish znf143 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Zebrafish znf143 Antibody (C-term) - Protein Information**Name** znf143**Synonyms** staf**Function** Transcriptional activator. Activates the gene for selenocysteine tRNA (tRNA^{Sec}). Binds to

the activator element (AE) motif of the selenocysteine tRNA gene promoter (By similarity).

Cellular Location


Nucleus.

Zebrafish znf143 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Zebrafish znf143 Antibody (C-term) - Images



Western blot analysis of lysate from zebrafish muscle tissue lysate, using Zebrafish znf143 Antibody (C-term) (Cat. #Azb18703c). Azb18703c was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

Zebrafish znf143 Antibody (C-term) - Background

Transcriptional activator. Activates the gene for selenocysteine tRNA (tRNA^{sec}). Binds to the activator element (AE) motif of the selenocysteine tRNA gene promoter (By similarity).

Zebrafish znf143 Antibody (C-term) - References

Myslinski E., et al. Gene 330:149-158(2004).
Howe K., et al. Nature 496:498-503(2013).