

(DANRE) stk10 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb19128c

Specification

(DANRE) stk10 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q7SY52
Reactivity	Zebrafish
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	113687
Antigen Region	342-376

(DANRE) stk10 Antibody (Center) - Additional Information

Gene ID 394108

Other Names

Serine/threonine-protein kinase 10, stk10

Target/Specificity

This DANRE stk10 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 342-376 amino acids from the Central region of DANRE stk10.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(DANRE) stk10 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) stk10 Antibody (Center) - Protein Information

Name stk10

Function May act as a polo kinase kinase by mediating phosphorylation of plk1.

Cellular Location


Cell membrane; Peripheral membrane protein

(DANRE) stk10 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

(DANRE) stk10 Antibody (Center) - Images



Anti-stk10 Antibody (Center) at 1:2000 dilution + ZF4 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 114 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

(DANRE) stk10 Antibody (Center) - Background

May act as a polo kinase kinase by mediating phosphorylation of plk1.