

Zebrafish stk10 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # Azb19128c

Specification

Zebrafish stk10 Antibody (Center) - Product Information

Application

Primary Accession

Reactivity

Host

Clonality

Isotype

Antigen Region

WB,E

075Y52

Zebrafish

Rabbit

polyclonal

Rabbit IgG

342-376

Zebrafish stk10 Antibody (Center) - Additional Information

Gene ID 394108

Other Names

Serine/threonine-protein kinase 10, stk10

Target/Specificity

This Zebrafish stk10 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 342-376 amino acids from the central region of zebrafish stk10.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Zebrafish stk10 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Zebrafish stk10 Antibody (Center) - Protein Information

Name stk10

Function May act as a polo kinase kinase by mediating phosphorylation of plk1.

Cellular Location




Cell membrane; Peripheral membrane protein

Zebrafish stk10 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Zebrafish stk10 Antibody (Center) - Images



Anti-stk10 Antibody (Center)at 1:2000 dilution + ZF4 whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 114 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Zebrafish stk10 Antibody (Center) - Background

May act as a polo kinase kinase by mediating phosphorylation of plk1.