

APOBEC3C Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP10252b**Specification**

APOBEC3C Antibody (C-term) Blocking peptide - Product InformationPrimary Accession
Other Accession[O9NRW3](#)
[NP_055323.2](#)**APOBEC3C Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 27350**Other Names**

DNA dC->dU-editing enzyme APOBEC-3C, A3C, 354-, APOBEC1-like, Phorbolin I, APOBEC3C, APOBEC1L, PBI

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

APOBEC3C Antibody (C-term) Blocking peptide - Protein Information**Name** APOBEC3C**Synonyms** APOBEC1L, PBI**Function**

DNA deaminase (cytidine deaminase) which acts as an inhibitor of retrovirus replication and retrotransposon mobility via deaminase- dependent and -independent mechanisms. After the penetration of retroviral nucleocapsids into target cells of infection and the initiation of reverse transcription, it can induce the conversion of cytosine to uracil in the minus-sense single-strand viral DNA, leading to G-to-A hypermutations in the subsequent plus-strand viral DNA. The resultant detrimental levels of mutations in the proviral genome, along with a deamination-independent mechanism that works prior to the proviral integration, together exert efficient antiretroviral effects in infected target cells. Selectively targets single-stranded DNA and does not deaminate double-stranded DNA or single- or double-stranded RNA. Exhibits antiviral activity against simian immunodeficiency virus (SIV), hepatitis B virus (HBV), herpes simplex virus 1 (HHV-1) and Epstein-Barr virus (EBV) and may inhibit the mobility of LTR and non- LTR retrotransposons. May also play a role in the epigenetic regulation of gene expression through the process of active DNA demethylation.

Cellular Location

Nucleus. Cytoplasm

Tissue Location

Expressed in spleen, testes, peripheral blood lymphocytes, heart, thymus, prostate and ovary

APOBEC3C Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

APOBEC3C Antibody (C-term) Blocking peptide - Images**APOBEC3C Antibody (C-term) Blocking peptide - Background**

This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or cell cycle control. [provided by RefSeq].

APOBEC3C Antibody (C-term) Blocking peptide - References

Albin, J.S., et al. Expert Rev Mol Med 12, E4 (2010) : Prochnow, C., et al. Sci. China, C, Life Sci. 52(10):893-902(2009) Lin, H., et al. Invest. Ophthalmol. Vis. Sci. 50(9):4436-4443(2009) Stauch, B., et al. Proc. Natl. Acad. Sci. U.S.A. 106(29):12079-12084(2009) Ross, S.R. PLoS Pathog. 5 (4), E1000347 (2009) :