

# SPIRE2 Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP10600c

#### **Specification**

# **SPIRE2 Antibody (Center) Blocking Peptide - Product Information**

Primary Accession <u>Q8WWL2</u> Other Accession <u>NP 115827.1</u>

# SPIRE2 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 84501

#### **Other Names**

Protein spire homolog 2, Spir-2, SPIRE2, KIAA1832, SPIR2

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

#### SPIRE2 Antibody (Center) Blocking Peptide - Protein Information

Name SPIRE2

Synonyms KIAA1832, SPIR2

#### **Function**

Acts as an actin nucleation factor, remains associated with the slow-growing pointed end of the new filament (PubMed:<a href="http://www.uniprot.org/citations/21620703" target="\_blank">21620703</a>). Involved in intracellular vesicle transport along actin fibers, providing a novel link between actin cytoskeleton dynamics and intracellular transport (By similarity). Required for asymmetric spindle positioning and asymmetric cell division during meiosis (PubMed:<a href="http://www.uniprot.org/citations/21620703" target="\_blank">21620703" target="\_blank">21620703</a>). Required for normal formation of the cleavage furrow and for polar body extrusion during female germ cell meiosis (PubMed:<a href="http://www.uniprot.org/citations/21620703" target="\_blank">21620703</a>/a>). Also acts in the nucleus: together with SPIRE1 and SPIRE2, promotes assembly of nuclear actin filaments in response to DNA damage in order to facilitate movement of chromatin and repair factors after DNA damage (PubMed:<a href="http://www.uniprot.org/citations/26287480" target=" blank">26287480</a>).

# **Cellular Location**



Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q8K1S6}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q8K1S6}. Cell membrane {ECO:0000250|UniProtKB:Q8K1S6}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q8K1S6}; Cytoplasmic side {ECO:0000250|UniProtKB:Q8K1S6}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q8K1S6}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q8K1S6}; Cytoplasmic side {ECO:0000250|UniProtKB:Q8K1S6}. Note=Detected at the cleavage furrow during asymmetric oocyte division and polar body extrusion {ECO:0000250|UniProtKB:Q8K1S6}

#### SPIRE2 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

SPIRE2 Antibody (Center) Blocking Peptide - Images

# SPIRE2 Antibody (Center) Blocking Peptide - Background

Acts as a actin nucleation factor, remains associated with the slow-growing pointed end of the new filament. Involved in vesicle transport processes providing a novel link between actin organization and intracellular transport (By similarity).

#### SPIRE2 Antibody (Center) Blocking Peptide - References

Pechlivanis, M., et al. J. Biol. Chem. 284(37):25324-25333(2009)