

NR0B2 Antibody (Center) Blocking peptide

Synthetic peptide Catalog # BP11075c

Specification

NR0B2 Antibody (Center) Blocking peptide - Product Information

Primary Accession

Q15466

NR0B2 Antibody (Center) Blocking peptide - Additional Information

Gene ID 8431

Other Names

Nuclear receptor subfamily 0 group B member 2, Orphan nuclear receptor SHP, Small heterodimer partner, NR0B2, SHP

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NR0B2 Antibody (Center) Blocking peptide - Protein Information

Name NR0B2

Synonyms SHP

Function

Transcriptional regulator that acts as a negative regulator of receptor-dependent signaling pathways (By similarity). Specifically inhibits transactivation of the nuclear receptor with which it interacts (By similarity). Inhibits transcriptional activity of NEUROD1 on E-box-containing promoter by interfering with the coactivation function of the p300/CBP-mediated transcription complex for NEUROD1 (PubMed:14752053 (a>). Essential component of the liver circadian clock which via its interaction with NR1D1 and RORG regulates NPAS2-mediated hepatic lipid metabolism (By similarity). Regulates the circadian expression of cytochrome P450 (CYP) enzymes (By similarity). Represses: NR5A2 and HNF4A to down-regulate CYP2C38, NFLI3 to up-regulate CYP2A5, BHLHE41/HNF1A axis to up-regulate CYP1A2, CYP2E1 and CYP3A11, and NR1D1 to up-regulate CYP2B10, CYP4A10 and CYP4A14 (By similarity).

Cellular Location

Nucleus. Cytoplasm Note=Colocalizes with NEUROD1 in the nucleus



Tissue Location

Liver. Low levels of expression were detected in heart and pancreas.

NR0B2 Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

NR0B2 Antibody (Center) Blocking peptide - Images

NR0B2 Antibody (Center) Blocking peptide - Background

The protein encoded by this gene is an unusual orphanreceptor that contains a putative ligand-binding domain but lacks aconventional DNA-binding domain. The gene product is a member ofthe nuclear hormone receptor family, a group of transcription factors regulated by small hydrophobic hormones, a subset of whichdo not have known ligands and are referred to as orphan nuclear receptors. The protein has been shown to interact with retinoid and thyroid hormone receptors, inhibiting their ligand-dependent ranscriptional activation. In addition, interaction with estrogenreceptors has been demonstrated, leading to inhibition of function. Studies suggest that the protein represses nuclear hormonereceptor-mediated transactivation via two separate steps:competition with coactivators and the direct effects of its transcriptional repressor function.

NR0B2 Antibody (Center) Blocking peptide - References

Zhou, T., et al. J. Biol. Chem. 285(32):24871-24881(2010)Roberts, K.E., et al. Gastroenterology 139(1):130-139(2010)Kim, K., et al. Biochem. J. 427(3):413-422(2010)Yang, Z., et al. J. Int. Med. Res. 38(1):142-149(2010)Ondracek, C.R., et al. J. Virol. 83(23):12535-12544(2009)