

ARRDC3 antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP11631b**Specification**

ARRDC3 antibody (C-term) Blocking peptide - Product InformationPrimary Accession [Q96B67](#)**ARRDC3 antibody (C-term) Blocking peptide - Additional Information**

Gene ID 57561

Other Names

Arrestin domain-containing protein 3, TBP-2-like inducible membrane protein, TLIMP, ARRDC3, KIAA1376

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ARRDC3 antibody (C-term) Blocking peptide - Protein Information

Name ARRDC3

Synonyms KIAA1376

Function

Adapter protein that plays a role in regulating cell-surface expression of adrenergic receptors and probably also other G protein- coupled receptors (PubMed:20559325, PubMed:21982743, PubMed:23208550). Plays a role in NEDD4-mediated ubiquitination and endocytosis of activated ADRB2 and subsequent ADRB2 degradation (PubMed:20559325, PubMed:23208550). May recruit NEDD4 to ADRB2 (PubMed:20559325). Alternatively, may function as adapter protein that does not play a major role in recruiting NEDD4 to ADRB2, but rather plays a role in targeting ADRB2 to endosomes (PubMed:23208550).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side Lysosome Endosome Early endosome. Note=Associated with plasma membrane, as well as with endosomes and lysosomes during endocytosis (PubMed:16269462, PubMed:23208550, PubMed:20559325)

Tissue Location

Highly expressed in skeletal muscle, placenta, kidney, lung, liver, blood, adrenal gland, lymph node, mammary gland, thyroid, and trachea (PubMed:16269462, PubMed:21982743). Very low levels in colon, thymus, spleen, small intestine, bladder and bone marrow (PubMed:16269462). Strong expression in differentiated adipocytes compared to preadipocytes (PubMed:16269462). Detected in omental fat and subcutaneous fat tissue (PubMed:21982743)

ARRDC3 antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ARRDC3 antibody (C-term) Blocking peptide - Images**ARRDC3 antibody (C-term) Blocking peptide - Background**

The arrestins are a family of proteins that are important for regulating signal transduction within cells. Arrestins are part of a conserved two step mechanism for regulating the activity of G-protein coupled receptors (GPCRs). In response to a stimulus, GPCRs activate a heterotrimeric G protein. In order to turn off this response, or adapt to a constant stimulus, activated receptors need to be silenced. The first step is phosphorylation by a class of serine/threonine kinases called G protein coupled receptor kinases (GRKs). This phosphorylation specifically marks the activated receptor for arrestin binding. Once arrestin is bound to the receptor it is unable to signal further. Recent research continues to expand the known actions of arrestins, which can bind to other classes of receptors and can directly activate signaling pathways on their own. Different arrestins (visual arrestin (or Arrestin 1), beta-arrestin 1 (or Arrestin 2) and beta-arrestin 2 (or Arrestin 3) can reduce the activity of their target GPCRs in several different ways.

ARRDC3 antibody (C-term) Blocking peptide - References

Davis, O.S., et al. Behav. Genet. 40(6):759-767(2010) Draheim, K.M., et al. Oncogene 29(36):5032-5047(2010) Nabhan, J.F., et al. EMBO Rep. 11(8):605-611(2010) Patwari, P., et al. J. Biol. Chem. 284(37):24996-25003(2009) Lamesch, P., et al. Genomics 89(3):307-315(2007)