

### SLC7A2 Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP11660b

### **Specification**

### SLC7A2 Antibody (C-term) Blocking peptide - Product Information

Primary Accession

P52569

# SLC7A2 Antibody (C-term) Blocking peptide - Additional Information

**Gene ID 6542** 

#### **Other Names**

Cationic amino acid transporter 2, CAT-2, CAT2, Low affinity cationic amino acid transporter 2, Solute carrier family 7 member 2, SLC7A2, ATRC2, CAT2

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### SLC7A2 Antibody (C-term) Blocking peptide - Protein Information

### Name SLC7A2 (HGNC:11060)

### **Function**

Functions as a permease involved in the transport of the cationic amino acids (L-arginine, L-lysine, L-ornithine and L- homoarginine); the affinity for its substrates differs between isoforms created by alternative splicing (PubMed:<a href="http://www.uniprot.org/citations/28684763" target="\_blank">28684763</a>, PubMed:<a href="http://www.uniprot.org/citations/9174363" target="\_blank">9174363</a>). May play a role in classical or alternative activation of macrophages via its role in arginine transport (By similarity).

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein

#### **Tissue Location**

Expressed at high levels in the skeletal muscle, placenta and ovary. Expressed at intermediate levels in the liver and pancreas and at low levels in the kidney and heart

# SLC7A2 Antibody (C-term) Blocking peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

SLC7A2 Antibody (C-term) Blocking peptide - Images

# SLC7A2 Antibody (C-term) Blocking peptide - Background

Low-affinity, high capacity permease involved in the transport of the cationic amino acids (arginine, lysine and ornithine). Plays a regulatory role in classical or alternative activation of macrophages (By similarity).

# SLC7A2 Antibody (C-term) Blocking peptide - References

Chaturvedi, R., et al. Gastroenterology 139(5):1686-1698(2010)Visigalli, R., et al. J. Mol. Cell. Cardiol. 49(2):260-270(2010)Yokoyama, K., et al. Nephron Clin Pract 115 (4), C237-C243 (2010): Jaeger, K., et al. Histochem. Cell Biol. 129(3):321-329(2008)Broer, S. Physiol. Rev. 88(1):249-286(2008)