

SSH3 Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP11697b**Specification**

SSH3 Antibody (C-term) Blocking peptide - Product Information

Primary Accession [Q8TE77](#)

SSH3 Antibody (C-term) Blocking peptide - Additional Information

Gene ID 54961

Other Names

Protein phosphatase Slingshot homolog 3, SSH-like protein 3, SSH-3L, hSSH-3L, SSH3, SSH3L

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SSH3 Antibody (C-term) Blocking peptide - Protein Information

Name SSH3

Synonyms SSH3L

Function

Protein phosphatase which may play a role in the regulation of actin filament dynamics. Can dephosphorylate and activate the actin binding/depolymerizing factor cofilin, which subsequently binds to actin filaments and stimulates their disassembly (By similarity).

Cellular Location

Cytoplasm, cytoskeleton. Nucleus

SSH3 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SSH3 Antibody (C-term) Blocking peptide - Images

SSH3 Antibody (C-term) Blocking peptide - Background

This gene is a member of the protocadherin beta genecluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor geneclusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections.

SSH3 Antibody (C-term) Blocking peptide - References

Zhang, Z., et al. Protein Sci. 13(10):2819-2824(2004) Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003) Frank, M., et al. Curr. Opin. Cell Biol. 14(5):557-562(2002) Vanhalst, K., et al. FEBS Lett. 495 (1-2), 120-125 (2001) :Wu, Q., et al. Genome Res. 11(3):389-404(2001)