

RPS6KB1 Antibody (T389) Blocking peptide
Synthetic peptide
Catalog # BP11736a**Specification**

RPS6KB1 Antibody (T389) Blocking peptide - Product Information

Primary Accession [P23443](#)
Other Accession [NP_003152.1](#)

RPS6KB1 Antibody (T389) Blocking peptide - Additional Information

Gene ID 6198

Other Names

Ribosomal protein S6 kinase beta-1, S6K-beta-1, S6K1, 70 kDa ribosomal protein S6 kinase 1, P70S6K1, p70-S6K 1, Ribosomal protein S6 kinase I, Serine/threonine-protein kinase 14A, p70 ribosomal S6 kinase alpha, p70 S6 kinase alpha, p70 S6K-alpha, p70 S6KA, RPS6KB1, STK14A

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

RPS6KB1 Antibody (T389) Blocking peptide - Protein Information

Name RPS6KB1

Synonyms STK14A

Function

Serine/threonine-protein kinase that acts downstream of mTOR signaling in response to growth factors and nutrients to promote cell proliferation, cell growth and cell cycle progression (PubMed:11500364, PubMed:12801526, PubMed:14673156, PubMed:15071500, PubMed:15341740, PubMed:16286006, PubMed:17052453, PubMed:17053147, PubMed:17936702, PubMed:18952604, PubMed:19085255,</p>
</div>
<div data-bbox="437 955 499 971" data-label="Page-Footer">
 Page 1/3
</div>

PubMed:19720745,
PubMed:19935711,
PubMed:19995915,
PubMed:23429703,
PubMed:28178239,
PubMed:22017876).
Regulates protein synthesis through phosphorylation of EIF4B, RPS6 and EEF2K, and contributes to cell survival by repressing the pro-apoptotic function of BAD (PubMed:11500364), PubMed:12801526), PubMed:14673156), PubMed:15071500), PubMed:15341740), PubMed:16286006), PubMed:17052453), PubMed:17053147), PubMed:17936702), PubMed:18952604), PubMed:19085255), PubMed:19720745), PubMed:19935711), PubMed:19995915), PubMed:23429703), PubMed:28178239), PubMed:22017876). Under conditions of nutrient depletion, the inactive form associates with the EIF3 translation initiation complex (PubMed:16286006). Upon mitogenic stimulation, phosphorylation by the mechanistic target of rapamycin complex 1 (mTORC1) leads to dissociation from the EIF3 complex and activation (PubMed:16286006). The active form then phosphorylates and activates several substrates in the pre-initiation complex, including the EIF2B complex and the cap-binding complex component EIF4B (PubMed:16286006). Also controls translation initiation by phosphorylating a negative regulator of EIF4A, PDCD4, targeting it for ubiquitination and subsequent proteolysis (PubMed:17053147). Promotes initiation of the pioneer round of protein synthesis by phosphorylating POLDIP3/SKAR (PubMed:15341740). In response to IGF1, activates translation elongation by phosphorylating EEF2 kinase (EEF2K), which leads to its inhibition and thus activation of EEF2 (PubMed:11500364). Also plays a role in feedback regulation of mTORC2 by mTORC1 by phosphorylating RICTOR, resulting in the inhibition of mTORC2 and AKT1 signaling (PubMed:19720745), PubMed:19935711), PubMed:19995915). Also involved in feedback regulation of mTORC1 and mTORC2 by phosphorylating DEPTOR (PubMed:22017876). Mediates cell survival by phosphorylating the pro- apoptotic protein BAD and suppressing its pro-apoptotic function (By similarity). Phosphorylates mitochondrial URI1 leading to dissociation of a URI1-PPP1CC complex (PubMed:17936702). The free mitochondrial PPP1CC can then dephosphorylate RPS6KB1 at Thr-412, which is proposed to be a negative feedback mechanism for the RPS6KB1 anti-apoptotic function (PubMed:17936702). Mediates TNF-alpha-induced insulin resistance by phosphorylating IRS1 at multiple serine residues, resulting in accelerated degradation of IRS1 (PubMed:18952604).

In cells lacking functional TSC1-2 complex, constitutively phosphorylates and inhibits GSK3B (PubMed:17052453). May be involved in cytoskeletal rearrangement through binding to neurabin (By similarity). Phosphorylates and activates the pyrimidine biosynthesis enzyme CAD, downstream of MTOR (PubMed:23429703). Following activation by mTORC1, phosphorylates EPRS and thereby plays a key role in fatty acid uptake by adipocytes and also most probably in interferon-gamma-induced translation inhibition (PubMed:28178239).

Cellular Location

Synapse, synaptosome. Mitochondrion outer membrane. Mitochondrion. Note=Colocalizes with URI1 at mitochondrion [Isoform Alpha II]: Cytoplasm.

Tissue Location

Widely expressed..

RPS6KB1 Antibody (T389) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

RPS6KB1 Antibody (T389) Blocking peptide - Images**RPS6KB1 Antibody (T389) Blocking peptide - Background**

This gene encodes a member of the RSK (ribosomal S6kinase) family of serine/threonine kinases. This kinase contains 2non-identical kinase catalytic domains and phosphorylates severalresidues of the S6 ribosomal protein. The kinase activity of thisprotein leads to an increase in protein synthesis and cellproliferation. Amplification of the region of DNA encoding thisgene and overexpression of this kinase are seen in some breastcancer cell lines. Alternate translational start sites have beendescribed and alternate transcriptional splice variants have beenobserved but have not been thoroughly characterized. [provided byRefSeq].

RPS6KB1 Antibody (T389) Blocking peptide - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)Lai, K.P., et al. EMBO J. 29(17):2994-3006(2010)Treekitkarnmongkol, W., et al. World J. Gastroenterol. 16(32):4047-4054(2010)Fenton, T.R., et al. Biochem. Biophys. Res. Commun. 398(3):400-405(2010)Bian, C.X., et al. Biochem. Biophys. Res. Commun. 398(3):395-399(2010)