

# KCNH6 Antibody (Center) Blocking peptide

Synthetic peptide Catalog # BP11929c

## **Specification**

## KCNH6 Antibody (Center) Blocking peptide - Product Information

Primary Accession

Q9H252

## KCNH6 Antibody (Center) Blocking peptide - Additional Information

**Gene ID** 81033

#### **Other Names**

Potassium voltage-gated channel subfamily H member 6, Ether-a-go-go-related gene potassium channel 2, ERG-2, Eag-related protein 2, Ether-a-go-go-related protein 2, hERG-2, hERG2, Voltage-gated potassium channel subunit Kv112, KCNH6, ERG2

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# KCNH6 Antibody (Center) Blocking peptide - Protein Information

Name KCNH6

**Synonyms** ERG2

## **Function**

Pore-forming (alpha) subunit of voltage-gated potassium channel. Elicits a slowly activating, rectifying current (By similarity). Channel properties may be modulated by cAMP and subunit assembly.

### **Cellular Location**

Membrane; Multi-pass membrane protein.

### **Tissue Location**

Expressed in prolactin-secreting adenomas.

#### KCNH6 Antibody (Center) Blocking peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

# KCNH6 Antibody (Center) Blocking peptide - Images

# KCNH6 Antibody (Center) Blocking peptide - Background

Voltage-gated potassium (Kv) channels represent the mostcomplex class of voltage-gated ion channels from both functionaland structural standpoints. Their diverse functions includeregulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smoothmuscle contraction, and cell volume. This gene encodes a member ofthe potassium channel, voltage-gated, subfamily H. This member is apore-forming (alpha) subunit. Several alternatively splicedtranscript variants have been identified from this gene, but thefull-length nature of only two transcript variants has beendetermined.

## KCNH6 Antibody (Center) Blocking peptide - References

Liu, B.C., et al. Prog. Neuropsychopharmacol. Biol. Psychiatry 34(3):506-509(2010)Gutman, G.A., et al. Pharmacol. Rev. 57(4):473-508(2005)Bauer, C.K., et al. Pflugers Arch. 445(5):589-600(2003)Ganetzky, B., et al. Ann. N. Y. Acad. Sci. 868, 356-369 (1999):