

**MED6 Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP12093b****Specification**

---

**MED6 Antibody (C-term) Blocking peptide - Product Information**Primary Accession [O75586](#)**MED6 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 10001**Other Names**

Mediator of RNA polymerase II transcription subunit 6, Activator-recruited cofactor 33 kDa component, ARC33, Mediator complex subunit 6, hMed6, Renal carcinoma antigen NY-REN-28, MED6, ARC33

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**MED6 Antibody (C-term) Blocking peptide - Protein Information****Name** MED6**Synonyms** ARC33**Function**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

**Cellular Location**

Nucleus.

**MED6 Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **MED6 Antibody (C-term) Blocking peptide - Images**

#### **MED6 Antibody (C-term) Blocking peptide - Background**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

#### **MED6 Antibody (C-term) Blocking peptide - References**

Pavri, R., et al. Mol. Cell 18(1):83-96(2005) Sato, S., et al. Mol. Cell 14(5):685-691(2004) Yang, F., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2339-2344(2004) Tomomori-Sato, C., et al. J. Biol. Chem. 279(7):5846-5851(2004) Sato, S., et al. J. Biol. Chem. 278(50):49671-49674(2003)