

LIN7C Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP12807b**Specification**

LIN7C Antibody (C-term) Blocking peptide - Product InformationPrimary Accession [Q9NUP9](#)**LIN7C Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 55327**Other Names**

Protein lin-7 homolog C, Lin-7C, Mammalian lin-seven protein 3, MALS-3, Vertebrate lin-7 homolog 3, Veli-3, LIN7C, MALS3, VELI3

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

LIN7C Antibody (C-term) Blocking peptide - Protein Information**Name** LIN7C**Synonyms** MALS3, VELI3**Function**

Plays a role in establishing and maintaining the asymmetric distribution of channels and receptors at the plasma membrane of polarized cells. Forms membrane-associated multiprotein complexes that may regulate delivery and recycling of proteins to the correct membrane domains. The tripartite complex composed of LIN7 (LIN7A, LIN7B or LIN7C), CASK and APBA1 associates with the motor protein KIF17 to transport vesicles containing N-methyl-D-aspartate (NMDA) receptor subunit NR2B along microtubules (By similarity). This complex may have the potential to couple synaptic vesicle exocytosis to cell adhesion in brain. Ensures the proper localization of GRIN2B (subunit 2B of the NMDA receptor) to neuronal postsynaptic density and may function in localizing synaptic vesicles at synapses where it is recruited by beta-catenin and cadherin. Required to localize Kir2 channels, GABA transporter (SLC6A12) and EGFR/ERBB1, ERBB2, ERBB3 and ERBB4 to the basolateral membrane of epithelial cells.

Cellular Location

Cell membrane; Peripheral membrane protein. Basolateral cell membrane; Peripheral membrane protein. Cell junction Postsynaptic density membrane; Peripheral membrane protein. Cell junction,

tight junction Note=Mainly basolateral in renal epithelial cells.

LIN7C Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

LIN7C Antibody (C-term) Blocking peptide - Images

LIN7C Antibody (C-term) Blocking peptide - Background

LIN7C plays a role in establishing and maintaining the asymmetric distribution of channels and receptors at the plasma membrane of polarized cells. Forms membrane-associated multiprotein complexes that may regulate delivery and recycling of proteins to the correct membrane domains. The tripartite complex composed of LIN7 (LIN7A, LIN7B or LIN7C), CASK and APBA1 may have the potential to couple synaptic vesicle exocytosis to cell adhesion in brain. Ensures the proper localization of GRIN2B (subunit 2B of the NMDA receptor) to neuronal postsynaptic density and may function in localizing synaptic vesicles at synapses where it is recruited by beta-catenin and cadherin. Required to localize Kir2 channels, GABA transporter (SLC6A12) and EGFR/ERBB1, ERBB2, ERBB3 and ERBB4 to the basolateral membrane of epithelial cells.

LIN7C Antibody (C-term) Blocking peptide - References

Ng, M.C., et al. J. Clin. Endocrinol. Metab. 95(5):2418-2425(2010)Lanktree, M., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 147B (6), 945-951 (2008) :Bulgin, N.L., et al. Neuromolecular Med. 10(4):343-355(2008)Onda, T., et al. Cancer Res. 67(20):9643-9648(2007)Bohl, J., et al. J. Biol. Chem. 282(13):9392-9400(2007)