

Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide
Synthetic peptide
Catalog # BP1303a**Specification**

Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide - Product InformationPrimary Accession [P10415](#)**Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide - Additional Information****Gene ID** 596**Other Names**

Apoptosis regulator Bcl-2, BCL2

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP1303a](/product/products/AP1303a) was selected from the region of human Bcl-2 BH3 Domain. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide - Protein Information**Name** BCL2**Function**

Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells (PubMed:[1508712](http://www.uniprot.org/citations/1508712), PubMed:[8183370](http://www.uniprot.org/citations/8183370)). Regulates cell death by controlling the mitochondrial membrane permeability (PubMed:[11368354](http://www.uniprot.org/citations/11368354)). Appears to function in a feedback loop system with caspases (PubMed:[11368354](http://www.uniprot.org/citations/11368354)). Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1) (PubMed:[11368354](http://www.uniprot.org/citations/11368354)). Also acts as an inhibitor of autophagy: interacts with BECN1 and AMBRA1 during non-starvation conditions and inhibits their autophagy function (PubMed:[18570871](http://www.uniprot.org/citations/18570871))

target="_blank">18570871, PubMed:20889974, PubMed:21358617). May attenuate inflammation by impairing NLRP1-inflammasome activation, hence CASP1 activation and IL1B release (PubMed:17418785).

Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein. Cytoplasm {ECO:0000250|UniProtKB:P10417}

Tissue Location

Expressed in a variety of tissues.

Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide - Images

Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide - Background

BCL2 is integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends.

Bcl-2 Antibody (BH3 Domain Specific) Blocking peptide - References

Zhu, C.J., et al., J. Neurosci. Res. 74(1):60-66 (2003).Scopa, C.D., et al., Dig. Dis. Sci. 48(10):1990-1997 (2003).Grace, V.M., et al., Gynecol. Oncol. 91(1):51-58 (2003).Thoulouze, M.I., et al., Virology 314(2):549-561 (2003).Iwata, A., et al., J. Immunol. 171(6):3136-3141 (2003).