

**UNC5D Antibody (Center) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP13072c****Specification**

---

**UNC5D Antibody (Center) Blocking peptide - Product Information**Primary Accession [Q6UXZ4](#)**UNC5D Antibody (Center) Blocking peptide - Additional Information****Gene ID** 137970**Other Names**

Netrin receptor UNC5D, Protein unc-5 homolog 4, Protein unc-5 homolog D, UNC5D, KIAA1777, UNC5H4

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**UNC5D Antibody (Center) Blocking peptide - Protein Information****Name** UNC5D**Synonyms** KIAA1777, UNC5H4 {ECO:0000303|Ref.2}**Function**

Receptor for the netrin NTN4 that promotes neuronal cell survival (By similarity). Plays a role in cell-cell adhesion and cell guidance. Receptor for netrin involved in cell migration. Plays a role in axon guidance by mediating axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding (By similarity). May play a role in apoptosis in response to DNA damage (PubMed:<a href="http://www.uniprot.org/citations/24691657" target="\_blank">24691657</a>). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (PubMed:<a href="http://www.uniprot.org/citations/24519068" target="\_blank">24519068</a>). Mediates cell-cell adhesion via its interaction with FLRT3 on an adjacent cell (By similarity).

**Cellular Location**

Cell membrane; Single-pass type I membrane protein

## **UNC5D Antibody (Center) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

## **UNC5D Antibody (Center) Blocking peptide - Images**

## **UNC5D Antibody (Center) Blocking peptide - Background**

UNC5D is receptor for netrin. May be involved in axon guidance by mediating axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (By similarity).

## **UNC5D Antibody (Center) Blocking peptide - References**

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :Wang, H., et al. Biochem. Biophys. Res. Commun. 370(4):594-598(2008)Choy, K.W., et al. Physiol. Genomics 25(1):9-15(2006)Zhong, Y., et al. Cereb. Cortex 14(10):1144-1152(2004)Beausoleil, S.A., et al. Proc. Natl. Acad. Sci. U.S.A. 101(33):12130-12135(2004)