

RPL34 Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP13207c

Specification

RPL34 Antibody (Center) Blocking Peptide - Product Information

Primary Accession

RPL34 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 6164

Other Names

60S ribosomal protein L34, RPL34

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13207c was selected from the Center region of RPL34. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

P49207

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

RPL34 Antibody (Center) Blocking Peptide - Protein Information

Name RPL34

Function

Component of the large ribosomal subunit (PubMed:12962325, PubMed:23636399, PubMed:25957688, PubMed:25901680, PubMed:32669547). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed:12962325, PubMed:23636399, PubMed:25957688, PubMed:25901680, PubMed:25901680, PubMed:25901680, PubMed:25901680, PubMed:32669547, PubMed:32669547.



Tel: 858.875.1900 Fax: 858.875.1999

Cellular Location

Cytoplasm, cytosol. Cytoplasm Endoplasmic reticulum {ECO:0000250|UniProtKB:Q29223}. Note=Detected on cytosolic polysomes (PubMed:25957688). Detected in ribosomes that are associated with the rough endoplasmic reticulum (By similarity) {ECO:0000250|UniProtKB:Q29223, ECO:0000269|PubMed:25957688}

RPL34 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

RPL34 Antibody (Center) Blocking Peptide - Images

RPL34 Antibody (Center) Blocking Peptide - Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Togetherthese subunits are composed of 4 RNA species and approximately 80structurally distinct proteins. This gene encodes a ribosomalprotein that is a component of the 60S subunit. The protein belongsto the L34E family of ribosomal proteins. It is located in thecytoplasm. This gene originally was thought to be located at 17g21, but it has been mapped to 4q. Transcript variants derived fromalternative splicing, alternative transcription initiation sites, and/or alternative polyadenylation exist; these variants encode thesame protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersedthrough the genome.

RPL34 Antibody (Center) Blocking Peptide - References

Kalsi, G., et al. Hum. Mol. Genet. 19(12):2497-2506(2010)Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :Andersen, J.S., et al. Nature 433(7021):77-83(2005)Kapp, L.D., et al. Annu. Rev. Biochem. 73, 657-704 (2004) :Mazumder, B., et al. Cell 115(2):187-198(2003)