

OR8I2 Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP13261b**Specification**

OR8I2 Antibody (C-term) Blocking peptide - Product InformationPrimary Accession [Q8NOY5](#)**OR8I2 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 120586**Other Names**

Olfactory receptor 8I2, Olfactory receptor OR11-170, OR8I2

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13261b was selected from the C-term region of OR8I2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

OR8I2 Antibody (C-term) Blocking peptide - Protein Information**Name** OR8I2**Function**

Odorant receptor.

Cellular Location

Cell membrane; Multi-pass membrane protein.

OR8I2 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

OR8I2 Antibody (C-term) Blocking peptide - Images

OR8I2 Antibody (C-term) Blocking peptide - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR8I2 Antibody (C-term) Blocking peptide - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) ; Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589 (2004) ; Fuchs, T., et al. Genomics 80(3):295-302 (2002)