

SLIT1 Antibody (N-term) Blocking peptide
Synthetic peptide
Catalog # BP13374a**Specification**

SLIT1 Antibody (N-term) Blocking peptide - Product Information

Primary Accession [O75093](#)

SLIT1 Antibody (N-term) Blocking peptide - Additional Information

Gene ID 6585

Other Names

Slit homolog 1 protein, Slit-1, Multiple epidermal growth factor-like domains protein 4, Multiple EGF-like domains protein 4, SLIT1, KIAA0813, MEGF4, SLIL1

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13374a was selected from the N-term region of SLIT1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SLIT1 Antibody (N-term) Blocking peptide - Protein Information

Name SLIT1

Synonyms KIAA0813, MEGF4, SLIL1

Function

Thought to act as molecular guidance cue in cellular migration, and function appears to be mediated by interaction with roundabout homolog receptors. During neural development involved in axonal navigation at the ventral midline of the neural tube and projection of axons to different regions (By similarity). SLIT1 and SLIT2 together seem to be essential for midline guidance in the forebrain by acting as repulsive signal preventing inappropriate midline crossing by axons projecting from the olfactory bulb.

Cellular Location

Secreted.

Tissue Location

Predominantly expressed in adult forebrain. Expressed in fetal brain, lung and kidney

SLIT1 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SLIT1 Antibody (N-term) Blocking peptide - Images**SLIT1 Antibody (N-term) Blocking peptide - Background**

Thought to act as molecular guidance cue in cellular migration, and function appears to be mediated by interaction with roundabout homolog receptors. During neural development involved in axonal navigation at the ventral midline of the neural tube and projection of axons to different regions (By similarity). SLIT1 and SLIT2 together seem to be essential for midline guidance in the forebrain by acting as repulsive signal preventing inappropriate midline crossing by axons projecting from the olfactory bulb.

SLIT1 Antibody (N-term) Blocking peptide - References

Marlow, R., et al. Cancer Res. 68(19):7819-7827(2008) Hussain, S.A., et al. J. Biol. Chem. 281(51):39693-39698(2006) Lin, L., et al. Stem Cells 24(11):2504-2513(2006) Lim, J., et al. Cell 125(4):801-814(2006) Grupe, A., et al. Am. J. Hum. Genet. 78(1):78-88(2006)