

DHX32 Antibody (Center) Blocking peptide
Synthetic peptide
Catalog # BP13575c**Specification**

DHX32 Antibody (Center) Blocking peptide - Product InformationPrimary Accession [Q7L7V1](#)**DHX32 Antibody (Center) Blocking peptide - Additional Information****Gene ID** 55760**Other Names**

Putative pre-mRNA-splicing factor ATP-dependent RNA helicase DHX32, DEAD/H box 32, DEAD/H helicase-like protein 1, DHLP1, DEAH box protein 32, HuDDX32, DHX32, DDX32

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13575c was selected from the Center region of DHX32. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DHX32 Antibody (Center) Blocking peptide - Protein Information**Name** DHX32**Synonyms** DDX32**Cellular Location**

Nucleus. Mitochondrion

Tissue Location

Expressed in lymphoid tissues (at protein level). Expressed in brain, heart, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung, lymphoid tissues and blood leukocytes.

DHX32 Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

DHX32 Antibody (Center) Blocking peptide - Images

DHX32 Antibody (Center) Blocking peptide - Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this DEAD box protein family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a member of this family. The function of this member has not been determined. Alternative splicing of this gene generates 2 transcript variants, but the full length nature of one of the variants has not been defined.

DHX32 Antibody (Center) Blocking peptide - References

Huang, C., et al. J. Exp. Clin. Cancer Res. 28, 11 (2009) :Chen, Y., et al. Exp. Mol. Pathol. 82(3):256-261(2007)Alli, Z., et al. Anticancer Res. 27 (1A), 373-377 (2007) :Alli, Z., et al. Exp. Mol. Pathol. 81(3):245-248(2006)Alli, Z., et al. Cell. Immunol. 237(2):141-146(2005)