

PPP3CA Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP13757b

Specification

PPP3CA Antibody (C-term) Blocking peptide - Product Information

Primary Accession

Q08209

PPP3CA Antibody (C-term) Blocking peptide - Additional Information

Gene ID 5530

Other Names

Serine/threonine-protein phosphatase 2B catalytic subunit alpha isoform, CAM-PRP catalytic subunit, Calmodulin-dependent calcineurin A subunit alpha isoform, PPP3CA, CALNA, CNA

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13757b was selected from the C-term region of PPP3CA. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PPP3CA Antibody (C-term) Blocking peptide - Protein Information

Name PPP3CA (HGNC:9314)

Synonyms CALNA, CNA

Function

Calcium-dependent, calmodulin-stimulated protein phosphatase which plays an essential role in the transduction of intracellular Ca(2+)-mediated signals (PubMed:15671020, PubMed:18838687, PubMed:19154138, PubMed:23468591, PubMed:23468591, PubMed:30254215). Many of the

substrates contain a PXIXIT motif and/or a LxVP motif (PubMed:17498738, PubMed:17502104, PubMed:<a



href="http://www.uniprot.org/citations/22343722" target=" blank">22343722, PubMed:23468591, PubMed:27974827). In response to increased Ca(2+) levels, dephosphorylates and activates phosphatase SSH1 which results in cofilin dephosphorylation (PubMed: 15671020). In response to increased Ca(2+) levels following mitochondrial depolarization, dephosphorylates DNM1L inducing DNM1L translocation to the mitochondrion (PubMed:18838687). Positively regulates the CACNA1B/CAV2.2-mediated Ca(2+) release probability at hippocampal neuronal soma and synaptic terminals (By similarity). Dephosphorylates heat shock protein HSPB1 (By similarity). Dephosphorylates and activates transcription factor NFATC1 (PubMed:19154138). In response to increased Ca(2+) levels, regulates NFAT-mediated transcription probably by dephosphorylating NFAT and promoting its nuclear translocation (PubMed:<a $href="http://www.uniprot.org/citations/26248042"\ target="_blank">26248042).$ Dephosphorylates and inactivates transcription factor ELK1 (PubMed:19154138). Dephosphorylates DARPP32 (PubMed:19154138). May dephosphorylate CRTC2 at 'Ser-171' resulting in CRTC2 dissociation from 14-3-3 proteins (PubMed: 30611118). Dephosphorylates transcription factor TFEB at 'Ser- 211' following Coxsackievirus B3 infection, promoting nuclear translocation (PubMed:33691586). Required for postnatal development of the nephrogenic zone and superficial glomeruli in the kidneys, cell cycle homeostasis in the nephrogenic zone, and ultimately normal kidney function (By similarity). Plays a role in intracellular AQP2 processing and localization to the apical membrane in the kidney, may thereby be required for efficient kidney filtration (By similarity). Required for secretion of salivary enzymes amylase, peroxidase, lysozyme and sialic acid via formation of secretory vesicles in the submandibular glands (By similarity). Required for calcineurin activity and homosynaptic depotentiation in the hippocampus (By similarity). Required for normal differentiation and survival of keratinocytes and therefore required for epidermis superstructure formation (By similarity). Positively regulates osteoblastic bone formation, via promotion of osteoblast differentiation (By similarity). Positively regulates osteoclast differentiation, potentially via NFATC1 signaling (By similarity). May play a role in skeletal muscle fiber type specification, potentially via NFATC1 signaling (By similarity). Negatively regulates MAP3K14/NIK signaling via inhibition of nuclear translocation of the transcription factors RELA and RELB (By similarity). Required for antigen-specific T- cell proliferation response (By similarity). Dephosphorylates KLHL3, promoting the interaction between KLHL3 and WNK4 and subsequent degradation of WNK4 (PubMed: 30718414). Negatively regulates SLC9A1 activity (PubMed: 31375679).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:P63329}. Cytoplasm, myofibril, sarcomere, Z line {ECO:0000250|UniProtKB:P63329}. Cell projection, dendritic spine. Note=Colocalizes with ACTN1 and MYOZ2 at the Z line in heart and skeletal muscle (By similarity). Recruited to the cell membrane by scaffold protein AKAP5 following L-type Ca(2+)-channel activation (PubMed:22343722) {ECO:0000250|UniProtKB:P63329, ECO:0000269|PubMed:22343722}

Tissue Location

Expressed in keratinocytes (at protein level) (PubMed:29043977). Expressed in lymphoblasts (at protein level) (PubMed:30254215).

PPP3CA Antibody (C-term) Blocking peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

PPP3CA Antibody (C-term) Blocking peptide - Images

PPP3CA Antibody (C-term) Blocking peptide - Background

Calcium-dependent, calmodulin-stimulated protein phosphatase. This subunit may have a role in the calmodulin activation of calcineurin. Dephosphorylates DNM1L, HSPB1 and SSH1.

PPP3CA Antibody (C-term) Blocking peptide - References

He, Z.H., et al. Eur. J. Appl. Physiol. 110(4):761-767(2010)Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)Chiocco, M.J., et al. Subst Use Misuse 45(11):1809-1826(2010)Yang, D., et al. Int. J. Mol. Med. 26(1):159-164(2010)Bollo, M., et al. PLoS ONE 5 (8), E11925 (2010):