

Clorf187 Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP13829b

Specification

Clorf187 Antibody (C-term) Blocking peptide - Product Information

Primary Accession

Q8NBI3

Clorf187 Antibody (C-term) Blocking peptide - Additional Information

Gene ID 374946

Other Names

Draxin {ECO:0000255|HAMAP-Rule:MF_03060}, Dorsal inhibitory axon guidance protein {ECO:0000255|HAMAP-Rule:MF_03060}, Dorsal repulsive axon guidance protein {ECO:0000255|HAMAP-Rule:MF_03060}, Neucrin, DRAXIN {ECO:0000255|HAMAP-Rule:MF_03060}, C1orf187

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13829b was selected from the C-term region of Clorf187. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Clorf187 Antibody (C-term) Blocking peptide - Protein Information

Name DRAXIN {ECO:0000255|HAMAP-Rule:MF 03060}

Synonyms Clorf187

Function

Chemorepulsive axon guidance protein required for the development of spinal cord and forebrain commissures. Acts as a chemorepulsive guidance protein for commissural axons during development. Able to inhibit or repel neurite outgrowth from dorsal spinal cord. Inhibits the stabilization of cytosolic beta-catenin (CTNNB1) via its interaction with LRP6, thereby acting as an antagonist of Wnt signaling pathway.

Cellular Location

Secreted.



Clorf187 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

Clorf187 Antibody (C-term) Blocking peptide - Images

Clorf187 Antibody (C-term) Blocking peptide - Background

Chemorepulsive axon guidance protein required for the development of spinal cord and forebrain commissures. Acts as a chemorepulsive guidance protein for commissural axons during development. Able to inhibit or repel neurite outgrowth from dorsal spinal cord. Inhibits the stabilization of cytosolic beta-catenin (CTNNB1) via its interaction with LRP6, thereby acting as an antagonist of Wnt signaling pathway (By similarity).

Clorf187 Antibody (C-term) Blocking peptide - References

Miyake, A., et al. Biochem. Biophys. Res. Commun. 390(3):1051-1055(2009)Islam, S.M., et al. Science 323(5912):388-393(2009)Sullivan, P.F., et al. Mol. Psychiatry 13(6):570-584(2008)Zhang, Z., et al. Protein Sci. 13(10):2819-2824(2004)Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003)