

**AGR2 Antibody (N-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP13845a**

### Specification

#### AGR2 Antibody (N-term) Blocking peptide - Product Information

Primary Accession [O95994](#)

#### AGR2 Antibody (N-term) Blocking peptide - Additional Information

##### Gene ID 10551

##### Other Names

Anterior gradient protein 2 homolog, AG-2, hAG-2, HPC8, Secreted cement gland protein XAG-2 homolog, AGR2, AG2

##### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13845a was selected from the N-term region of AGR2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

##### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

##### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

##### Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

#### AGR2 Antibody (N-term) Blocking peptide - Protein Information

##### Name AGR2

##### Synonyms AG2

##### Function

Required for MUC2 post-transcriptional synthesis and secretion. May play a role in the production of mucus by intestinal cells (By similarity). Proto-oncogene that may play a role in cell migration, cell differentiation and cell growth. Promotes cell adhesion (PubMed:<a href="http://www.uniprot.org/citations/23274113" target="\_blank">23274113</a>).

##### Cellular Location

Secreted. Endoplasmic reticulum {ECO:0000250|UniProtKB:O88312}

##### Tissue Location

Expressed strongly in trachea, lung, stomach, colon, prostate and small intestine. Expressed

weakly in pituitary gland, salivary gland, mammary gland, bladder, appendix, ovary, fetal lung, uterus, pancreas, kidney, fetal kidney, testis, placenta, thyroid gland and in estrogen receptor (ER)-positive breast cancer cell lines

### **AGR2 Antibody (N-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **AGR2 Antibody (N-term) Blocking peptide - Images**

### **AGR2 Antibody (N-term) Blocking peptide - Background**

Required for MUC2 post-transcriptional synthesis and secretion. May play a role in the production of mucus by intestinal cells (By similarity). Proto-oncogene that may play a role in cell migration, cell differentiation and cell growth.

### **AGR2 Antibody (N-term) Blocking peptide - References**

Ambolet-Camoit, A., et al. *Toxicol. Sci.* 115(2):501-512(2010) Lipkin, S.M., et al. *Cancer Prev Res (Phila)* 3(5):597-603(2010) Chen, R., et al. *Mol. Cancer* 9, 149 (2010) :Vanderlaag, K.E., et al. *Breast Cancer Res.* 12 (3), R32 (2010) :Rice, G.E., et al. *J. Exp. Clin. Cancer Res.* 29, 62 (2010) :