

# ADAMTS5 Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP13847a

# **Specification**

# ADAMTS5 Antibody (N-term) Blocking peptide - Product Information

**Primary Accession** 

**Q9UNA0** 

# ADAMTS5 Antibody (N-term) Blocking peptide - Additional Information

**Gene ID** 11096

#### **Other Names**

A disintegrin and metalloproteinase with thrombospondin motifs 5, ADAM-TS 5, ADAM-TS5, ADAMTS-5, 3424-, A disintegrin and metalloproteinase with thrombospondin motifs 11, ADAM-TS 11, ADAMTS-11, ADMP-2, Aggrecanase-2, ADAMTS5, ADAMTS11, ADMP2

### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13847a was selected from the N-term region of ADAMTS5. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# ADAMTS5 Antibody (N-term) Blocking peptide - Protein Information

Name ADAMTS5

Synonyms ADAMTS11, ADMP2

### **Function**

Metalloproteinase that plays an important role in connective tissue organization, development, inflammation and cell migration. Extracellular matrix (ECM) degrading enzyme that show proteolytic activity toward the hyalectan group of chondroitin sulfate proteoglycans (CSPGs) including ACAN, VCAN, BCAN and NCAN (PubMed:<a

href="http://www.uniprot.org/citations/16133547" target="\_blank">16133547</a>, PubMed:<a href="http://www.uniprot.org/citations/18992360" target="\_blank">18992360</a>). Cleavage within the hyalectans occurs at Glu-Xaa recognition motifs. Plays a role in embryonic development, including limb and cardiac morphogenesis, and skeletal muscle development through its VCAN remodeling properties. Cleaves VCAN in the pericellular matrix surrounding myoblasts, facilitating



Tel: 858.875.1900 Fax: 858.875.1999

myoblast contact and fusion which is required for skeletal muscle development and regeneration (By similarity). Participates in development of brown adipose tissue and browning of white adipose tissue (By similarity). Plays an important role for T-lymphocyte migration from draining lymph nodes following viral infection.

### **Cellular Location**

Secreted, extracellular space, extracellular matrix

### **Tissue Location**

Expressed at low level in placenta primarily but also detected in heart and brain, cervix, uterus, bladder, esophagus, rib cartilage, chondroblastoma, fibrous tissue and a joint capsule from an arthritic patient

# ADAMTS5 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

# Blocking Peptides

ADAMTS5 Antibody (N-term) Blocking peptide - Images

# ADAMTS5 Antibody (N-term) Blocking peptide - Background

ADAMTS5 cleaves aggrecan, a cartilage proteoglycan, and may be involved in its turnover. May play an important role in the destruction of aggrecan in arthritic diseases. May play a role in proteolytic processing mostly during the peri-implantation period.