

# MGMT Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP13848a

## **Specification**

# MGMT Antibody (N-term) Blocking peptide - Product Information

Primary Accession P16455

# MGMT Antibody (N-term) Blocking peptide - Additional Information

**Gene ID 4255** 

#### **Other Names**

Methylated-DNA--protein-cysteine methyltransferase, 6-O-methylguanine-DNA methyltransferase, MGMT, O-6-methylguanine-DNA-alkyltransferase, MGMT

# **Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP13848a was selected from the N-term region of MGMT. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

## **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# MGMT Antibody (N-term) Blocking peptide - Protein Information

**Name MGMT** 

## **Function**

Involved in the cellular defense against the biological effects of O6-methylguanine (O6-MeG) and O4-methylthymine (O4-MeT) in DNA. Repairs the methylated nucleobase in DNA by stoichiometrically transferring the methyl group to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.

## **Cellular Location**

Nucleus.

### MGMT Antibody (N-term) Blocking peptide - Protocols





Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

MGMT Antibody (N-term) Blocking peptide - Images

MGMT Antibody (N-term) Blocking peptide - Background

MGMT is involved in the cellular defense against the biological effects of O6-methylguanine (O6-MeG) in DNA. Repairs alkylated guanine in DNA by stoichiometrically transferring the alkyl group at the O-6 position to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.