

NRTN Antibody (Center) Blocking peptide

Synthetic peptide Catalog # BP14055c

Specification

NRTN Antibody (Center) Blocking peptide - Product Information

Primary Accession

Q99748

NRTN Antibody (Center) Blocking peptide - Additional Information

Gene ID 4902

Other Names Neurturin, NRTN

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP14055c was selected from the Center region of NRTN. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NRTN Antibody (Center) Blocking peptide - Protein Information

Name NRTN

Function

Supports the survival of sympathetic neurons in culture. May regulate the development and maintenance of the CNS. Might control the size of non-neuronal cell population such as haemopoietic cells.

Cellular Location

Secreted.

NRTN Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.



• Blocking Peptides

NRTN Antibody (Center) Blocking peptide - Images

NRTN Antibody (Center) Blocking peptide - Background

Neurturin is a member of the TGF-beta subfamily, TRN. Thisgene signals through RET and a GPI-linked coreceptor, and promotessurvival of neuronal populations. A neurturin mutation has been described in a family with Hirschsprung Disease. [provided byRefSeq].

NRTN Antibody (Center) Blocking peptide - References

Fernandez, R.M., et al. J. Pediatr. Surg. 43(11):2042-2047(2008)Adly, M.A., et al. J. Am. Acad. Dermatol. 58(2):238-250(2008)Yoong, L.F., et al. J. Neurosci. 27(21):5603-5614(2007)Yoong, L.F., et al. J. Neurochem. 98(4):1149-1158(2006)Vargas-Leal, V., et al. J. Immunol. 175(4):2301-2308(2005)