

Mouse Pkmyt1 Antibody (N-term) Blocking peptide
Synthetic peptide
Catalog # BP14078a**Specification**

Mouse Pkmyt1 Antibody (N-term) Blocking peptide - Product InformationPrimary Accession [Q9ESG9](#)**Mouse Pkmyt1 Antibody (N-term) Blocking peptide - Additional Information****Gene ID** 268930**Other Names**

Membrane-associated tyrosine- and threonine-specific cdc2-inhibitory kinase, Myt1 kinase, Pkmyt1, Myt1

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP14078a was selected from the N-term region of Mouse Pkmyt1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Mouse Pkmyt1 Antibody (N-term) Blocking peptide - Protein Information**Name** Pkmyt1**Synonyms** Myt1**Function**

Acts as a negative regulator of entry into mitosis (G2 to M transition) by phosphorylation of the CDK1 kinase specifically when CDK1 is complexed to cyclins. Mediates phosphorylation of CDK1 predominantly on 'Thr-14'. Also involved in Golgi fragmentation. May be involved in phosphorylation of CDK1 on 'Tyr-15' to a lesser degree, however tyrosine kinase activity is unclear and may be indirect.

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q99640}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q99640}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q99640}; Peripheral membrane protein

{ECO:0000250|UniProtKB:Q99640}

Mouse Pkmyt1 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Mouse Pkmyt1 Antibody (N-term) Blocking peptide - Images

Mouse Pkmyt1 Antibody (N-term) Blocking peptide - Background

Pkmyt1 acts as a negative regulator of entry into mitosis (G2 to M transition) by phosphorylation of the CDK1 kinase specifically when CDK1 is complexed to cyclins. Mediates phosphorylation of CDK1 predominantly on 'Thr-14'. Also involved in Golgi fragmentation. May be involved in phosphorylation of CDK1 on 'Tyr-15' to a lesser degree, however tyrosine kinase activity is unclear and may be indirect. May be a downstream target of Notch signaling pathway during eye development (By similarity).