

# **CEBPE Antibody (C-term) Blocking peptide**

Synthetic peptide Catalog # BP14112b

## **Specification**

# **CEBPE Antibody (C-term) Blocking peptide - Product Information**

**Primary Accession** 

**Q15744** 

# CEBPE Antibody (C-term) Blocking peptide - Additional Information

**Gene ID 1053** 

#### **Other Names**

CCAAT/enhancer-binding protein epsilon, C/EBP epsilon, CEBPE

### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP14112b was selected from the C-term region of CEBPE. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# **CEBPE Antibody (C-term) Blocking peptide - Protein Information**

## Name CEBPE

#### **Function**

Transcriptional activator (PubMed:<a href="http://www.uniprot.org/citations/26019275" target="\_blank">26019275</a>). C/EBP are DNA- binding proteins that recognize two different motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers. Required for the promyelocyte-myelocyte transition in myeloid differentiation (PubMed:<a href="http://www.uniprot.org/citations/10359588" target="blank">10359588</a>).

### **Cellular Location**

**Nucleus** 

# **Tissue Location**

Strongest expression occurs in promyelocyte and late-myeloblast-like cell lines.



# CEBPE Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

### Blocking Peptides

CEBPE Antibody (C-term) Blocking peptide - Images

# CEBPE Antibody (C-term) Blocking peptide - Background

The protein encoded by this gene is a bZIP transcriptionfactor which can bind as a homodimer to certain DNA regulatoryregions. It can also form heterodimers with the related proteinCEBP-delta. The encoded protein may be essential for terminal differentiation and functional maturation of committed granulocyteprogenitor cells. Mutations in this gene have been associated with Specific Granule Deficiency, a rare congenital disorder. Multiplevariants of this gene have been described, but the full-lengthnature of only one has been determined.

### **CEBPE Antibody (C-term) Blocking peptide - References**

Prasad, R.B., et al. Blood (2009) In press :Papaemmanuil, E., et al. Nat. Genet. 41(9):1006-1010(2009)Bedi, R., et al. Blood 113(2):317-327(2009)Cloutier, A., et al. J. Immunol. 182(1):563-571(2009)Matsushita, H., et al. Oncogene 27(53):6749-6760(2008)