

### **HNMT Antibody (C-term) Blocking Peptide**

Synthetic peptide Catalog # BP1442b

### **Specification**

### **HNMT Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession

P50135

## HNMT Antibody (C-term) Blocking Peptide - Additional Information

**Gene ID 3176** 

#### **Other Names**

Histamine N-methyltransferase, HMT, HNMT

#### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP1442b>AP1442b</a> was selected from the C-term region of human HNMT. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **HNMT Antibody (C-term) Blocking Peptide - Protein Information**

### **Name HNMT**

#### **Function**

Inactivates histamine by N-methylation. Plays an important role in degrading histamine and in regulating the airway response to histamine.

### **Cellular Location**

Cytoplasm.

## **HNMT Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.



# • Blocking Peptides

#### HNMT Antibody (C-term) Blocking Peptide - Images

# **HNMT Antibody (C-term) Blocking Peptide - Background**

In mammals, histamine is metabolized by two major pathways: N(tau)-methylation via histamine N-methyltransferase and oxidative deamination via diamine oxidase. This gene encodes the first enzyme which is found in the cytosol and uses S-adenosyl-L-methionine as the methyl donor. In the mammalianbrain, the neurotransmitter activity of histamine is controlled by N(tau)-methylation as diamine oxidase is not found in the central nervous system. A common genetic polymorphism affects the activity levels of this gene product in red blood cells.

### **HNMT Antibody (C-term) Blocking Peptide - References**

Aksoy, S., et al., Biochem. Biophys. Res. Commun. 219(2):548-554 (1996).