

DPF2 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP1451a**Specification**

DPF2 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q92785](#)**DPF2 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 5977**Other Names**

Zinc finger protein ubi-d4, Apoptosis response zinc finger protein, BRG1-associated factor 45D, BAF45D, D4, zinc and double PHD fingers family 2, Protein requiem, DPF2, BAF45D, REQ, UBID4

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP1451a](/product/products/AP1451a) was selected from the N-term region of human DPF2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DPF2 Antibody (N-term) Blocking Peptide - Protein Information**Name** DPF2**Synonyms** BAF45D, REQ, UBID4**Function**

Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed: [28533407](http://www.uniprot.org/citations/28533407), PubMed: [27775714](http://www.uniprot.org/citations/27775714)). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed: [28533407](http://www.uniprot.org/citations/28533407)). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B pathway (PubMed: [20460684](http://www.uniprot.org/citations/20460684)).

Cellular Location

Nucleus. Cytoplasm

Tissue Location

Ubiquitous.

DPF2 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

DPF2 Antibody (N-term) Blocking Peptide - Images**DPF2 Antibody (N-term) Blocking Peptide - Background**

DPF2 is a member of the d4 domain family, characterized by a zinc finger-like structural motif. This protein functions as a transcription factor which is necessary for the apoptotic response following deprivation of survival factors. It likely serves a regulatory role in rapid hematopoietic cell growth and turnover. DPF2 gene is considered a causal candidate for multiple endocrine neoplasia type I, an inherited cancer syndrome involving multiple parathyroid, enteropancreatic, and pituitary tumors.

DPF2 Antibody (N-term) Blocking Peptide - References

Olsen,J.V.,Cell 127 (3), 635-648 (2006)Beausoleil,S.A.,Proc. Natl. Acad. Sci. U.S.A. 101 (33), 12130-12135 (2004)