

AP2M1 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP14642b**Specification**

AP2M1 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [O96CW1](#)**AP2M1 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 1173

Other Names

AP-2 complex subunit mu, AP-2 mu chain, Adaptin-mu2, Adaptor protein complex AP-2 subunit mu, Adaptor-related protein complex 2 subunit mu, Clathrin assembly protein complex 2 mu medium chain, Clathrin coat assembly protein AP50, Clathrin coat-associated protein AP50, HA2 50 kDa subunit, Plasma membrane adaptor AP-2 50 kDa protein, AP2M1, CLAPM1, KIAA0109

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

AP2M1 Antibody (C-term) Blocking Peptide - Protein InformationName AP2M1 ([HGNC:564](#))

Synonyms CLAPM1, KIAA0109

Function

Component of the adaptor protein complex 2 (AP-2) (PubMed: [12694563](http://www.uniprot.org/citations/12694563), PubMed: [12952941](http://www.uniprot.org/citations/12952941), PubMed: [14745134](http://www.uniprot.org/citations/14745134), PubMed: [14985334](http://www.uniprot.org/citations/14985334), PubMed: [15473838](http://www.uniprot.org/citations/15473838), PubMed: [31104773](http://www.uniprot.org/citations/31104773)). Adaptor protein complexes function in protein transport via transport vesicles in different membrane traffic pathways (PubMed: [12694563](http://www.uniprot.org/citations/12694563), PubMed: [12952941](http://www.uniprot.org/citations/12952941), PubMed: [14745134](http://www.uniprot.org/citations/14745134), PubMed: [14985334](http://www.uniprot.org/citations/14985334), PubMed: [15473838](http://www.uniprot.org/citations/15473838)).

target="_blank">15473838, PubMed:31104773). Adaptor protein complexes are vesicle coat components and appear to be involved in cargo selection and vesicle formation (PubMed:12694563, PubMed:12952941, PubMed:14745134, PubMed:14985334, PubMed:15473838, PubMed:31104773). AP-2 is involved in clathrin-dependent endocytosis in which cargo proteins are incorporated into vesicles surrounded by clathrin (clathrin-coated vesicles, CCVs) which are destined for fusion with the early endosome (PubMed:12694563, PubMed:12952941, PubMed:14745134, PubMed:14985334, PubMed:15473838, PubMed:31104773). The clathrin lattice serves as a mechanical scaffold but is itself unable to bind directly to membrane components (PubMed:12694563, PubMed:12952941, PubMed:14745134, PubMed:14985334, PubMed:15473838, PubMed:31104773). Clathrin-associated adaptor protein (AP) complexes which can bind directly to both the clathrin lattice and to the lipid and protein components of membranes are considered to be the major clathrin adaptors contributing the CCV formation (PubMed:12694563, PubMed:12952941, PubMed:14745134, PubMed:14985334, PubMed:15473838, PubMed:31104773). AP-2 also serves as a cargo receptor to selectively sort the membrane proteins involved in receptor-mediated endocytosis (PubMed:16581796). AP-2 seems to play a role in the recycling of synaptic vesicle membranes from the presynaptic surface (PubMed:12694563, PubMed:12952941, PubMed:14745134, PubMed:14985334, PubMed:15473838, PubMed:31104773). AP-2 recognizes Y-X-X-[FILMV] (Y-X- X-Phi) and [ED]-X-X-X-L-[LI] endocytosis signal motifs within the cytosolic tails of transmembrane cargo molecules (By similarity). AP-2 may also play a role in maintaining normal post-endocytic trafficking through the ARF6-regulated, non-clathrin pathway (PubMed:19033387). During long-term potentiation in hippocampal neurons, AP-2 is responsible for the endocytosis of ADAM10 (PubMed:23676497). The AP-2 mu subunit binds to transmembrane cargo proteins; it recognizes the Y- X-X-Phi motifs (By similarity). The surface region interacting with to the Y-X-X-Phi motif is inaccessible in cytosolic AP-2, but becomes accessible through a conformational change following phosphorylation of AP-2 mu subunit at Thr-156 in membrane-associated AP-2 (PubMed:11877457). The membrane-specific phosphorylation event appears to involve assembled clathrin which activates the AP-2 mu kinase AAK1 (PubMed:11877457).

target="_blank">11877457). Plays a role in endocytosis of frizzled family members upon Wnt signaling (By similarity).

Cellular Location

Cell membrane. Membrane, coated pit; Peripheral membrane protein; Cytoplasmic side.
Note=AP-2 appears to be excluded from internalizing CCVs and to disengage from sites of endocytosis seconds before internalization of the nascent CCV {ECO:0000250|UniProtKB:P84091}

Tissue Location

Expressed in the brain (at protein level).

AP2M1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

AP2M1 Antibody (C-term) Blocking Peptide - Images

AP2M1 Antibody (C-term) Blocking Peptide - Background

This gene encodes a subunit of the heterotetrameric coat assembly protein complex 2 (AP2), which belongs to the adaptor complexes medium subunits family. The encoded protein is required for the activity of a vacuolar ATPase, which is responsible for proton pumping occurring in the acidification of endosomes and lysosomes. The encoded protein may also play an important role in regulating the intracellular trafficking and function of CTLA-4 protein. Two transcript variants encoding different isoforms have been found for this gene.

AP2M1 Antibody (C-term) Blocking Peptide - References

Kahlfeldt, N., et al. J. Biol. Chem. 285(4):2734-2749(2010) Boucrot, E., et al. PLoS ONE 5 (5), E10597 (2010) :Yoshida, T., et al. Int. J. Mol. Med. 24(2):233-246(2009) Levecque, C., et al. Am. J. Physiol., Cell Physiol. 297 (1), C160-C168 (2009) :Stanasila, L., et al. Mol. Pharmacol. 74(3):562-573(2008)