

ALDH3A2 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP1468b**Specification****ALDH3A2 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [P51648](#)**ALDH3A2 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID 224****Other Names**

Fatty aldehyde dehydrogenase, Aldehyde dehydrogenase 10, Aldehyde dehydrogenase family 3 member A2, Microsomal aldehyde dehydrogenase, ALDH3A2, ALDH10, FALDH

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP1468b was selected from the C-term region of human ALDH3A2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ALDH3A2 Antibody (C-term) Blocking Peptide - Protein Information**Name ALDH3A2****Function**

Catalyzes the oxidation of medium and long chain aliphatic aldehydes to fatty acids. Active on a variety of saturated and unsaturated aliphatic aldehydes between 6 and 24 carbons in length (PubMed:18035827, PubMed:18182499, PubMed:22633490, PubMed:25047030, PubMed:9133646, PubMed:9662422). Responsible for conversion of the sphingosine 1-phosphate (S1P) degradation product hexadecenal to hexadecenoic acid (PubMed:22633490).

Cellular Location

Microsome membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein; Cytoplasmic side {ECO:0000250|UniProtKB:P30839}

Tissue Location

Detected in liver (at protein level).

ALDH3A2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ALDH3A2 Antibody (C-term) Blocking Peptide - Images**ALDH3A2 Antibody (C-term) Blocking Peptide - Background**

Aldehyde dehydrogenase isozymes are thought to play a major role in the detoxification of aldehydes generated by alcohol metabolism and lipid peroxidation. ALDH3A2 catalyzes the oxidation of long-chain aliphatic aldehydes to fatty acid. Mutations in the ALDH3A2 gene cause Sjogren-Larsson syndrome.

ALDH3A2 Antibody (C-term) Blocking Peptide - References

Ashibe,B., J. Biol. Chem. 282 (28), 20763-20773 (2007)Rizzo,W.B., Mol. Genet. Metab. 90 (1), 1-9 (2007)