

**ALDH6A1 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP1469b****Specification**

---

**ALDH6A1 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q02252](#)**ALDH6A1 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 4329**Other Names**

Methylmalonate-semialdehyde dehydrogenase [acylating], mitochondrial, MMSDH,  
Malonate-semialdehyde dehydrogenase [acylating], Aldehyde dehydrogenase family 6 member  
A1, ALDH6A1, MMSDH

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP1469b](/product/products/AP1469b) was selected from the C-term region of human ALDH6A1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**ALDH6A1 Antibody (C-term) Blocking Peptide - Protein Information****Name** ALDH6A1 ([HGNC:7179](#))**Function**

Malonate and methylmalonate semialdehyde dehydrogenase involved in the catabolism of valine, thymine, and compounds catabolized by way of beta-alanine, including uracil and cytidine.

**Cellular Location**

Mitochondrion.

**ALDH6A1 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **ALDH6A1 Antibody (C-term) Blocking Peptide - Images**

#### **ALDH6A1 Antibody (C-term) Blocking Peptide - Background**

ALDH6A1 belongs to the aldehyde dehydrogenases family of proteins. This enzyme plays a role in the valine and pyrimidine catabolic pathways. This protein is a mitochondrial methylmalonate semialdehyde dehydrogenase, and catalyzes the irreversible oxidative decarboxylation of malonate and methylmalonate semialdehydes to acetyl- and propionyl-CoA. Methylmalonate semialdehyde dehydrogenase deficiency is characterized by elevated beta-alanine, 3-hydroxypropionic acid, and both isomers of 3-amino and 3-hydroxyisobutyric acids in urine organic acids.

#### **ALDH6A1 Antibody (C-term) Blocking Peptide - References**

Kuiper,H., Cytogenet. Genome Res. 109 (4), 533 (2005)Anderson,N.L., Mol. Cell Proteomics 3 (4), 311-326 (2004)Chambliss,K.L., J. Inherit. Metab. Dis. 23 (5), 497-504 (2000)Kedishvili,N.Y., J. Biol. Chem. 267 (27), 19724-19729 (1992)